

=> fil cap  
 FILE 'CAPLUS' ENTERED AT 12:12:19 ON 20 FEB 2008  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.


FILE COVERS 1907 - 20 Feb 2008 VOL 148 ISS 8  
 FILE LAST UPDATED: 19 Feb 2008 (20080219/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> d que 132

L1 STR



VAR G1=H/X  
 VAR G2=H/14/15  
 VAR G3=18/22  
 VAR G4=H/24  
 VAR G5=25/26/28/32/35  
 VAR G7=CHO/COOH  
 VAR G8=41/43/46  
 NODE ATTRIBUTES:  
 CONNECT IS E1 RC AT 14  
 CONNECT IS E2 RC AT 15  
 CONNECT IS E1 RC AT 24  
 CONNECT IS E1 RC AT 25  
 CONNECT IS E1 RC AT 27

CONNECT IS E2 RC AT 28  
 CONNECT IS E1 RC AT 34  
 CONNECT IS E1 RC AT 38  
 CONNECT IS E1 RC AT 42  
 CONNECT IS E1 RC AT 45  
 CONNECT IS E1 RC AT 46  
 DEFAULT MLEVEL IS ATOM  
 GGCAT IS LIN LOC SAT AT 14  
 GGCAT IS LIN LOC SAT AT 15  
 GGCAT IS LOC AT 24  
 GGCAT IS MCY UNS AT 25  
 GGCAT IS LOC AT 27  
 GGCAT IS MCY UNS AT 30  
 GGCAT IS MCY UNS AT 32  
 GGCAT IS MCY UNS AT 35  
 GGCAT IS SAT AT 45  
 GGCAT IS SAT AT 46  
 DEFAULT ECLEVEL IS LIMITED  
 ECOUNT IS E5 C E1 N AT 25  
 ECOUNT IS E6 C AT 30  
 ECOUNT IS E6 C AT 32  
 ECOUNT IS E6 C AT 35  
 ECOUNT IS M9 C AT 46

## GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED  
 NUMBER OF NODES IS 46

## STEREO ATTRIBUTES: NONE


L3 20 SEA FILE=REGISTRY SSS FUL L1  
 L30 1 SEA FILE=REGISTRY ABB=ON PLU=ON "BENZALDEHYDE, 3-CHLORO-6-HYD  
 ROXY-4-METHOXY-2-METHYL-5-((2E,6E)-3-METHYL-7-((2S)-TETRAHYDRO-  
 5,5-DIMETHYL-4-OXO-2-FURANYL)-2,6-OCTADIENYL)-"/CN  
 L31 21 SEA FILE=REGISTRY ABB=ON PLU=ON L3 OR L30  
 L32 3 SEA FILE=CAPLUS ABB=ON PLU=ON L31

=> d l32 ibib abs hitstr tot

L32 ANSWER 1 OF 3 CAPLUS COPYRIGHT 2008 ACS on STN  
 ACCESSION NUMBER: 2005:371202 CAPLUS Full-text  
 DOCUMENT NUMBER: 142:430014  
 TITLE: Preparation of phenol derivatives as anti-trypanosoma  
 agents  
 INVENTOR(S): Saimoto, Hiroyuki; Shigemasa, Yoshihiro; Kita,  
 Kiyoshi; Yabu, Yoshisada; Hosokawa, Tomoyoshi;  
 Yamamoto, Masaichi  
 PATENT ASSIGNEE(S): Arigen, Inc., Japan  
 SOURCE: PCT Int. Appl., 40 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2005037759	A1	20050428	WO 2003-JP13310	20031017
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,			
	CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,			

GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
 LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ,  
 OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,  
 TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,  
 KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,  
 FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR,  
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG  
 AU 2003273040 A1 20050505 AU 2003-273040 20031017  
 AU 2004282055 A1 20050428 AU 2004-282055 20041018  
 WO 2005037760 A1 20050428 WO 2004-JP15390 20041018  
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,  
 CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,  
 GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
 LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI,  
 NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,  
 TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW  
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,  
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,  
 EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE,  
 SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,  
 SN, TD, TG  
 EP 1681280 A1 20060719 EP 2004-792559 20041018  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK  
 CN 1882523 A 20061220 CN 2004-80033945 20041018  
 IN 2006DN02774 A 20070803 IN 2006-DN2774 20060517  
 US 2007208078 A1 20070906 US 2006-575653 20061213  
 PRIORITY APPLN. INFO.: WO 2003-JP13310 A 20031017  
 WO 2003-JP313310 A 20031017  
 WO 2004-JP15390 W 20041018  
 OTHER SOURCE(S): MARPAT 142:430014  
 GI



AB The title compds. I [X represents hydrogen or halogeno; R1 represents hydrogen, etc.; R2 represents hydrogen or C1-4 alkyl; R3 represents CHO or COOH; and R4 represents (CH<sub>2</sub>)<sub>m</sub>CH<sub>3</sub> (wherein m is an integer of 1 to 14), etc.] are prepared Thus, 2,4-dihydroxy-3-(1-hydroxydodecyl)-6-methylbenzaldehyde was prepared from 2,4-dihydroxy-6-methylbenzaldehyde and dodecanal. Compds. of this invention in vitro showed IC<sub>50</sub> values of 0.3 nM to 120 nM in an anti-trypanosoma assay.

IT 850732-56-8P 850732-57-9P 850732-58-0P  
 850732-59-1P 850732-60-4P 850732-61-5P  
 850732-62-6P 850732-63-7P 850732-64-8P  
 850732-65-9P 850732-66-0P 850732-67-1P


850732-68-2P 850732-69-3P 850732-70-6P  
 850732-71-7P 850732-72-8P 850732-73-9P  
 850732-74-0P 850732-75-1P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU  
 (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES  
 (Uses)

(preparation of phenol derivs. as anti-trypanosoma agents)


RN 850732-56-8 CAPLUS

CN Benzaldehyde, 2,4-dihydroxy-3-(1-hydroxydodecyl)-6-methyl- (CA INDEX  
 NAME)




RN 850732-57-9 CAPLUS

CN Benzaldehyde, 2,4-dihydroxy-3-(1-hydroxypropyl)-6-methyl- (CA INDEX NAME)




RN 850732-58-0 CAPLUS

CN Benzaldehyde, 2,4-dihydroxy-3-(1-hydroxypentyl)-6-methyl- (CA INDEX NAME)




RN 850732-59-1 CAPLUS

CN Benzaldehyde, 2,4-dihydroxy-3-(1-hydroxyheptyl)-6-methyl- (CA INDEX NAME)




RN 850732-60-4 CAPLUS

CN Benzaldehyde, 2,4-dihydroxy-3-(1-hydroxynonyl)-6-methyl- (CA INDEX NAME)




RN 850732-61-5 CAPLUS

CN Benzaldehyde, 2,4-dihydroxy-3-(1-hydroxydecyl)-6-methyl- (CA INDEX NAME)




RN 850732-62-6 CAPLUS

CN Benzaldehyde, 3-chloro-4,6-dihydroxy-5-(1-hydroxydodecyl)-2-methyl- (CA INDEX NAME)




RN 850732-63-7 CAPLUS


CN Benzaldehyde, 3-chloro-4,6-dihydroxy-5-(1-hydroxypropyl)-2-methyl- (CA INDEX NAME)




RN 850732-64-8 CAPLUS  
 CN Benzaldehyde, 3-chloro-4,6-dihydroxy-5-(1-hydroxypentyl)-2-methyl- (CA  
 INDEX NAME)




RN 850732-65-9 CAPLUS  
 CN Benzaldehyde, 3-chloro-4,6-dihydroxy-5-(1-hydroxyheptyl)-2-methyl- (CA  
 INDEX NAME)



RN 850732-66-0 CAPLUS  
 CN Benzaldehyde, 3-chloro-4,6-dihydroxy-5-(1-hydroxynonyl)-2-methyl- (CA  
 INDEX NAME)




RN 850732-67-1 CAPLUS  
 CN Benzaldehyde, 3-chloro-4,6-dihydroxy-5-(1-hydroxydecyl)-2-methyl- (CA  
 INDEX NAME)



RN 850732-68-2 CAPLUS

CN Benzaldehyde, 3-chloro-5-(1E)-1-dodecenyl-4,6-dihydroxy-2-methyl- (9CI)  
(CA INDEX NAME)


Double bond geometry as shown.



RN 850732-69-3 CAPLUS

CN Benzaldehyde, 3-(1E)-1-decenyl-2,4-dihydroxy- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 850732-70-6 CAPLUS

CN Benzaldehyde, 3-(1E)-1-dodecenyl-2,4-dihydroxy-6-methyl- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 850732-71-7 CAPLUS

CN Benzaldehyde, 3-chloro-4,6-dihydroxy-2-methyl-5-(1E)-1-propenyl- (9CI)  
(CA INDEX NAME)


Double bond geometry as shown.



RN 850732-72-8 CAPLUS

CN Benzaldehyde, 3-chloro-4,6-dihydroxy-2-methyl-5-(1E)-1-pentenyl- (9CI)  
(CA INDEX NAME)


Double bond geometry as shown.



RN 850732-73-9 CAPLUS

CN Benzaldehyde, 3-chloro-5-(1E)-1-heptenyl-4,6-dihydroxy-2-methyl- (9CI)  
(CA INDEX NAME)


Double bond geometry as shown.



RN 850732-74-0 CAPLUS

CN Benzaldehyde, 3-chloro-4,6-dihydroxy-2-methyl-5-(1E)-1-nonenyl- (9CI) (CA  
INDEX NAME)


Double bond geometry as shown.





RN 850732-75-1 CAPLUS  
 CN Benzaldehyde, 3-chloro-5-(1E)-1-decenyl-4,6-dihydroxy-2-methyl- (9CI) (CA  
 INDEX NAME)

Double bond geometry as shown.



REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L32 ANSWER 2 OF 3 CAPLUS COPYRIGHT 2008 ACS on STN  
 ACCESSION NUMBER: 2005:362051 CAPLUS Full-text  
 DOCUMENT NUMBER: 142:423803  
 TITLE: Prophylactic and therapeutic agents for  
 cryptosporidiosis containing ascochlorins,  
 ascofuranones, or dehydroascofuranones  
 INVENTOR(S): Kita, Kiyoshi; Yabu, Yoshisada; Nagai, Kazuo;  
 Minagawa, Nobuko; Hosokawa, Tomoyoshi; Suzuki,  
 Takashi; Ota, Nobuo  
 PATENT ASSIGNEE(S): Arigen, Inc., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 24 pp.  
 CODEN: JKXXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2005112755	A	20050428	JP 2003-347395	20031006
PRIORITY APPLN. INFO.:			JP 2003-347395	20031006
OTHER SOURCE(S):	MARPAT 142:423803			

GI

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB The agents contain  $\geq 1$  selected from ascochlorins I [ $R_1 = \text{CHO}, \text{CO}_2\text{H}$ ;  $R_2 = \text{H}$ ,  $(\text{C}_n\text{H}_{2n})\text{R}'$  ( $n = 1-5$ ;  $\text{R}' = \text{H}, \text{CO}_2\text{R}''$  on any C atom of  $\text{C}_n\text{H}_{2n}$ ;  $\text{R}'' = \text{H}, \text{C}_1-3$  alkyl),  $\text{COR}'''$  [ $\text{R}''' = \text{pyridyl}, \text{C}_1-3$  alkylamino, (halophenoxy)alkyl,  $\text{C}_1-3$  alkyl-Ph,  $(\text{C}_1-3$  alkoxy carbonyl)phenyl]], ascofuranones II, and dehydroascofuranones III, which inhibit cyanide-resistant quinol oxidase of *Cryptosporidium*. Thus,  $\text{IC}_{50}$  of ascofuranone against *Cryptosporidium* recombinant quinol oxidase was 0.3 nM. Tablets containing ascofuranone were also also formulated.

IT 611217-45-9  
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL  
 (Biological study); USES (Uses)


(prophylactic and therapeutic agents for cryptosporidiosis containing  
 ascochlorins, ascofuranones, or dehydroascofuranones as  
 cyanide-resistant quinol oxidase inhibitors)

RN 611217-45-9 CAPLUS

CN Benzaldehyde, 3-chloro-6-hydroxy-4-methoxy-2-methyl-5-[(2E,6E)-3-methyl-7-  
 [(2S)-tetrahydro-5,5-dimethyl-4-oxo-2-furanyl]-2,6-octadienyl]- (9CI) (CA  
 INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.



L32 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:643613 CAPLUS [Full-text](#)

DOCUMENT NUMBER: 139:307902

TITLE: Ascochlorin Derivatives as Ligands for Nuclear Hormone Receptors

AUTHOR(S): Togashi, Marie; Ozawa, Satoshi; Abe, Shoko; Nishimura, Tomoyuki; Tsuruga, Mie; Ando, Kunio; Tamura, Gakuzo; Kuwahara, Shigefumi; Ubukata, Makoto; Magae, Junji

CORPORATE SOURCE: Department of Biotechnology, Institute of Research and Innovation, Kashiwa, 277-0861, Japan

SOURCE: Journal of Medicinal Chemistry (2003), 46(19), 4113-4123

CODEN: JMCMAR; ISSN: 0022-2623


PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal


LANGUAGE: English

OTHER SOURCE(S): CASREACT 139:307902

GI



I



II

AB Nuclear receptor family proteins are structurally related transcription factors activated by specific lipophilic compds. Because they are activated by a variety of hormonal mols., including retinoic acid, vitamin D, and steroid hormones, they are assumed to be promising targets for clin. drugs. We previously found that one ascochlorin derivative, 4-O-carboxymethyl-ascochlorin, is a potent agonist of peroxisome proliferator activated receptor  $\gamma$  (PPAR $\gamma$ ). Here, we synthesized derivs. of ascochlorin, designated as a lead compound, to create new modulators of nuclear hormone receptors. Two derivs., 4-O-carboxymethyl-2-O-methylascochlorin and 4-O-isonicotinoyl-2-O-methylascochlorin, showed improved agonistic activity for PPAR $\gamma$  and induced differentiation of a progenitor cell line, C3H10T1/2. We also found that ascochlorin, dehydroascofuranone (I), and an ascochlorin 2,4-O-diacetyl-1-carboxylic acid derivative (II) specifically activated estrogen receptors, PPAR $\alpha$ , and an androgen receptor. All of the derivs. activated the pregnane X receptor. These results suggest that the chemical structure of ascochlorin is useful in designing novel modulators of nuclear receptors.

IT 611217-45-9P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(preparation of ascochlorin and ascofuranone derivs. as ligands for nuclear hormone receptors)

RN 611217-45-9 CAPLUS

CN Benzaldehyde, 3-chloro-6-hydroxy-4-methoxy-2-methyl-5-[(2E,6E)-3-methyl-7-[(2S)-tetrahydro-5,5-dimethyl-4-oxo-2-furanyl]-2,6-octadienyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.



REFERENCE COUNT: 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d his

(FILE 'HOME' ENTERED AT 11:10:38 ON 20 FEB 2008)

FILE 'REGISTRY' ENTERED AT 11:10:44 ON 20 FEB 2008

L1 STR  
L2 0 S L1  
L3 20 S L1 FUL  
L4 STR  
L5 3 S L4  
L6 46 S L4 FUL

FILE 'CAPLUS' ENTERED AT 11:39:00 ON 20 FEB 2008

L7 1 S L3  
L8 96 S L6

10/575,653

February 20, 2008

FILE 'REGISTRY' ENTERED AT 11:39:18 ON 20 FEB 2008

L9 STR L4  
L10 2 S L9 SAM SUB=L6  
L11 32 S L9 FUL SUB=L6

FILE 'CAPLUS' ENTERED AT 11:42:36 ON 20 FEB 2008

L12 93 S L11  
L13 ANALYZE L12 1-93 RN : 1785 TERMS

FILE 'REGISTRY' ENTERED AT 11:43:04 ON 20 FEB 2008

L14 1 S 38462-04-3  
L15 31 S L11 NOT L14

FILE 'CAPLUS' ENTERED AT 11:44:59 ON 20 FEB 2008

L16 34 S L15

FILE 'REGISTRY' ENTERED AT 11:45:09 ON 20 FEB 2008

L17 0 S L15 AND C18H25CLO4/MF  
L18 5 S L15 AND C=18  
L19 2 S L18 AND H=25  
L20 1 S L18 AND H=27  
L21 0 S L11 AND C=21  
L22 15 S L11 AND OC4/ES  
L23 15 S L22 AND O=5  
L24 0 S L23 AND N=1  
L25 0 S H=23 AND L23  
L26 3 S H=33 AND L23  
L27 3 S H=31 AND L23  
L28 0 S L26 AND C=23  
L29 3 S L11 AND C=24  
E BENZALDEHYDE, 3-CHLORO-6-HYDROXY-4-METHOXY-2-METHYL-5-((2E,6E  
L30 1 S E3  
L31 21 S L3 OR L30

FILE 'CAPLUS' ENTERED AT 12:03:19 ON 20 FEB 2008

L32 3 S L31

FILE 'CAPLUS, DISSABS, CONFSCI, WPIX' ENTERED AT 12:05:30 ON 20 FEB 2008

E SAIMOTO H/AU  
L33 206 S E3,E6-7  
E SHIGEMASA Y/AU  
L34 286 S E3,E5-6  
E KITA K/AU  
L35 1742 S E3-10,E80  
E YOSHISADA Y/AU  
E YABU Y/AU  
L36 85 S E3-4,E12  
E HOSOKAWA T/AU  
L37 1970 S E81,E3-11  
E YAMAMOTO M/AU  
L38 17371 S E140,E3-75  
L39 21463 S L33-38  
L40 286 S L39 AND PHENOL  
L41 0 S L40 AND ?PANASOM?  
L42 0 S L40 AND (TRYPANASOM? OR ?PANASOM? OR ANTITRYPANA?)

FILE 'CAPLUS' ENTERED AT 12:10:37 ON 20 FEB 2008

E US2006-575653/APPS  
L43 1 S E3

10/575,653

February 20, 2008

FILE 'CAPLUS, DISSABS, CONFSCI, WPIX' ENTERED AT 12:11:33 ON 20 FEB 2008  
L44 46 S L39 AND (?PANOSOM? OR TRYPANOS? OR ANTITRYPANOS?)

FILE 'CAPLUS' ENTERED AT 12:12:19 ON 20 FEB 2008

=> fil cap dissabs confsci wpix  
FILE 'CAPLUS' ENTERED AT 12:13:05 ON 20 FEB 2008  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'DISSABS' ENTERED AT 12:13:05 ON 20 FEB 2008  
COPYRIGHT (C) 2008 ProQuest Information and Learning Company; All Rights Reserved.

FILE 'CONFSCI' ENTERED AT 12:13:05 ON 20 FEB 2008  
COPYRIGHT (C) 2008 Cambridge Scientific Abstracts (CSA)

FILE 'WPIX' ENTERED AT 12:13:05 ON 20 FEB 2008  
COPYRIGHT (C) 2008 THE THOMSON CORPORATION

=> d que 144

L33 206 SEA ("SAIMOTO H"/AU OR "SAIMOTO HIROYUKI"/AU OR "SAIMOTO  
HIRYUKI"/AU)  
L34 286 SEA ("SHIGEMASA Y"/AU OR "SHIGEMASA YOSHIHIRO"/AU OR "SHIGEMASA  
YOSHIHRO"/AU)  
L35 1742 SEA ("KITA K"/AU OR "KITA K A"/AU OR "KITA K F"/AU OR "KITA K  
K F S P"/AU OR "KITA K M B"/AU OR "KITA K M C"/AU OR "KITA K N  
G K K K K"/AU OR "KITA K S C"/AU OR "KITA KIYOSHI"/AU)  
L36 85 SEA ("YABU Y"/AU OR "YABU Y T"/AU OR "YABU YOSHISADA"/AU)  
L37 1970 SEA ("HOSOKAWA TOMOYOSHI"/AU OR "HOSOKAWA T"/AU OR "HOSOKAWA T  
C O F"/AU OR "HOSOKAWA T D C"/AU OR "HOSOKAWA T F I C L"/AU OR  
"HOSOKAWA T I G C L"/AU OR "HOSOKAWA T L"/AU OR "HOSOKAWA T N  
D C"/AU OR "HOSOKAWA T T G C L"/AU OR "HOSOKAWA T Y F L"/AU)  
L38 17371 SEA ("YAMAMOTO MASAICHI"/AU OR "YAMAMOTO M"/AU OR "YAMAMOTO M  
O"/AU OR "YAMAMOTO M A"/AU OR "YAMAMOTO M B"/AU OR "YAMAMOTO M  
B K K K"/AU OR "YAMAMOTO M C"/AU OR "YAMAMOTO M C O M"/AU OR  
"YAMAMOTO M D"/AU OR "YAMAMOTO M D I L"/AU OR "YAMAMOTO M D K  
K"/AU OR "YAMAMOTO M D M"/AU OR "YAMAMOTO M D M C L"/AU OR  
"YAMAMOTO M D N P C L"/AU OR "YAMAMOTO M D P C C L"/AU OR  
"YAMAMOTO M D R L"/AU OR "YAMAMOTO M E"/AU OR "YAMAMOTO M E  
C"/AU OR "YAMAMOTO M E I"/AU OR "YAMAMOTO M EMILIA"/AU OR  
"YAMAMOTO M F"/AU OR "YAMAMOTO M F L"/AU OR "YAMAMOTO M F P F  
C L"/AU OR "YAMAMOTO M G C"/AU OR "YAMAMOTO M H I"/AU OR  
"YAMAMOTO M H L"/AU OR "YAMAMOTO M H M C L"/AU OR "YAMAMOTO M  
I"/AU OR "YAMAMOTO M I F D K K"/AU OR "YAMAMOTO M I P D N D  
I"/AU OR "YAMAMOTO M J"/AU OR "YAMAMOTO M J L"/AU OR "YAMAMOTO  
M K"/AU OR "YAMAMOTO M K C I C L"/AU OR "YAMAMOTO M K F"/AU OR  
"YAMAMOTO M K F S P"/AU OR "YAMAMOTO M K S S"/AU OR "YAMAMOTO  
M L"/AU OR "YAMAMOTO M M"/AU OR "YAMAMOTO M M C"/AU OR  
"YAMAMOTO M M C C"/AU OR "YAMAMOTO M M D K K"/AU OR "YAMAMOTO  
M M E W L"/AU OR "YAMAMOTO M M H I L"/AU OR "YAMAMOTO M M  
M"/AU OR "YAMAMOTO M M S K L"/AU OR "YAMAMOTO M M S K L P R  
C"/AU OR "YAMAMOTO M N"/AU OR "YAMAMOTO M N C I L"/AU OR  
"YAMAMOTO M N C N F"/AU OR "YAMAMOTO M N D C"/AU OR "YAMAMOTO  
M N D I"/AU OR "YAMAMOTO M N P C L"/AU OR "YAMAMOTO M O P C  
L"/AU OR "YAMAMOTO M O T"/AU OR "YAMAMOTO M P"/AU OR "YAMAMOTO  
M P I C L"/AU OR "YAMAMOTO M P R C"/AU OR "YAMAMOTO M P R C  
M"/AU OR "YAMAMOTO M R L"/AU OR "YAMAMOTO M S"/AU OR "YAMAMOTO  
M S C"/AU OR "YAMAMOTO M S E I L"/AU OR "YAMAMOTO M S K C  
L"/AU OR "YAMAMOTO M S L"/AU OR "YAMAMOTO M S L O G L"/AU OR

"YAMAMOTO M S M I L"/AU OR "YAMAMOTO M S P C L"/AU OR "YAMAMOTO  
M S S C C L"/AU OR "YAMAMOTO M T"/AU OR "YAMAMOTO M T L"/AU  
OR "YAMAMOTO M T L T K"/AU OR "YAMAMOTO M Y"/AU OR "YAMAMOTO M  
Y F C L"/AU)

L39 21463 SEA (L33 OR L34 OR L35 OR L36 OR L37 OR L38)  
L44 46 SEA L39 AND (?PANOSOM? OR TRYPANOS? OR ANTITRYPANOS?)

=> dup rem l44  
PROCESSING COMPLETED FOR L44  
L45 42 DUP REM L44 (4 DUPLICATES REMOVED)  
ANSWERS '1-40' FROM FILE CAPLUS  
ANSWER '41' FROM FILE CONFSCI  
ANSWER '42' FROM FILE WPIX

=> d l45 ibib abs tot

L45 ANSWER 1 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN DUPLICATE 1  
ACCESSION NUMBER: 2006:469037 CAPLUS Full-text  
DOCUMENT NUMBER: 144:482220  
TITLE: cDNA and protein sequences of Trypanosoma cruzi and  
Leishmania major quinol oxidase and screening its  
inhibitors for treatment of leishmaniasis and Chagas'  
diseases  
INVENTOR(S): Suzuki, Takashi; Suzuki, Mitsuko; Yabu, Yoshisada;  
Ota, Nobuo; Saimoto, Hiroyuki; Kita, Kiyoshi  
PATENT ASSIGNEE(S): Arigen, Inc., Japan  
SOURCE: Jpn. Kokai Tokkyo Koho, 25 pp.  
CODEN: JKXXAF  
DOCUMENT TYPE: Patent  
LANGUAGE: Japanese  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2006122010	A	20060518	JP 2004-317308	20041029
PRIORITY APPLN. INFO.:			JP 2004-317308	20041029

AB This invention provides cDNA and protein sequences of quinol oxidase from Trypanosoma cruzi and Leishmania major. This invention also provides Leishmania major quinol oxidase without cross reacting with AOX, a quinol oxidase from T. brucei. The enzyme activity of quinol oxidase was inhibited by ascofuranone and, the ascofuranone also inhibited the growth of Trypanosoma cruzi. Combining with cytochrome respiratory chain inhibitor, the inhibitors of quinol oxidase can be used for treatment of leishmaniasis and Chagas' diseases caused by infection of Trypanosoma cruzi and Leishmania major.

L45 ANSWER 2 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN DUPLICATE 2  
ACCESSION NUMBER: 2005:371202 CAPLUS Full-text  
DOCUMENT NUMBER: 142:430014  
TITLE: Preparation of phenol derivatives as  
anti-trypanosoma agents  
INVENTOR(S): Saimoto, Hiroyuki; Shigemasa, Yoshihiro; Kita,  
Kiyoshi; Yabu, Yoshisada; Hosokawa, Tomoyoshi;  
Yamamoto, Masaichi  
PATENT ASSIGNEE(S): Arigen, Inc., Japan  
SOURCE: PCT Int. Appl., 40 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent

LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005037759	A1	20050428	WO 2003-JP13310	20031017
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2003273040	A1	20050505	AU 2003-273040	20031017
AU 2004282055	A1	20050428	AU 2004-282055	20041018
WO 2005037760	A1	20050428	WO 2004-JP15390	20041018
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1681280	A1	20060719	EP 2004-792559	20041018
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK			
CN 1882523	A	20061220	CN 2004-80033945	20041018
IN 2006DN02774	A	20070803	IN 2006-DN2774	20060517
US 2007208078	A1	20070906	US 2006-575653	20061213
PRIORITY APPLN. INFO.:			WO 2003-JP13310	A 20031017
			WO 2003-JP313310	A 20031017
			WO 2004-JP15390	W 20041018
OTHER SOURCE(S):	MARPAT 142:430014			
GI				



AB The title compds. I [X represents hydrogen or halogeno; R1 represents hydrogen, etc.; R2 represents hydrogen or C1-4 alkyl; R3 represents CHO or COOH; and R4 represents (CH2)*m*CH3 (wherein *m* is an integer of 1 to 14), etc.] are prepared Thus, 2,4-dihydroxy-3-(1-hydroxydodecyl)-6- methylbenzaldehyde

was prepared from 2,4-dihydroxy-6-methylbenzaldehyde and dodecanal. Compds. of this invention in vitro showed IC50 values of 0.3 nM to 120 nM in an anti-trypanosoma assay.

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 3 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN DUPLICATE 3

ACCESSION NUMBER: 2004:677762 CAPLUS Full-text

DOCUMENT NUMBER: 141:167741

TITLE: Indole alkaloids as enhancers for antiprotozoal activity of ascofuranone, their compositions and kits, and treatment of protozoan diseases with them

INVENTOR(S): Kita, Kiyoshi; Yabu, Yoshisada; Nagai, Kazuo; Minagawa, Nobuko; Hosokawa, Kazuyoshi

PATENT ASSIGNEE(S): Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 21 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2004231601	A	20040819	JP 2003-24643	20030131
PRIORITY APPLN. INFO.:			JP 2003-24643	20030131
AB Title enhancers, useful for treatment of African trypanosoma, etc., contain indole alkaloids, e.g. in Picrasma quassioides. Thus, benzalharman at 25 $\mu$ M inhibited glycerokinase by 62.1%. Benzalharman enhanced the antiprotozoal activity of ascofuranone with ED50 of 8.5 $\mu$ M.				

L45 ANSWER 4 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN DUPLICATE 4

ACCESSION NUMBER: 1997:453440 CAPLUS Full-text

DOCUMENT NUMBER: 127:156717

TITLE: Protozoacides containing isoprenoid antibiotics, ascochlorin, ascofuranone, or their derivatives

INVENTOR(S): Minagawa, Nobuko; Yabu, Yoshisada; Kita, Kiyoshi; Nagai, Kazuo; Hosokawa, Tomoyoshi

PATENT ASSIGNEE(S): Hosokawa, Tomoyoshi, Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 6 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 09165332	A	19970624	JP 1995-351093	19951215
PRIORITY APPLN. INFO.:			JP 1995-351093	19951215
OTHER SOURCE(S):	MARPAT 127:156717			

GI





AB The protozoacides contain ascochlorins I [A = Q; R<sub>1</sub> = CHO, CO<sub>2</sub>H; R<sub>2</sub> = (C<sub>n</sub>H<sub>2n</sub>)R<sub>3</sub> (n = 1-5; R<sub>3</sub> = H, CO<sub>2</sub>R<sub>4</sub>; R<sub>4</sub> = H, C<sub>1</sub>-3 alkyl), COR<sub>5</sub> (R<sub>5</sub> = pyridyl, C<sub>1</sub>-3 alkylamino, halophenoxyalkyl, Ph substituted with C<sub>1</sub>-3 alkoxy or C<sub>1</sub>-3 alkoxy carbonyl)] or ascofuranones I (A = Q<sup>1</sup>) as active ingredients. The protozoacides are useful for prevention and treatment of African trypanosomiasis and trypanosomiasis of domestic animals. Ascochlorin inhibited in vitro growth of circulating-form *Trypanosoma brucei brucei* in the presence of glycerin. .

L45 ANSWER 5 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2007:1003374 CAPLUS Full-text

TITLE: *Trypanosoma brucei* vacuolar protein sorting 41 (VPS41) is required for intracellular iron utilization and maintenance of normal cellular morphology  
AUTHOR(S): Lu, S.; Suzuki, T.; Iizuka, N.; Ohshima, S.; Yabu, Y.; Suzuki, M.; Wen, L.; Ohta, N.

CORPORATE SOURCE: Department of Molecular Parasitology, Graduate School of Medical Sciences, Nagoya City University, Nagoya, 467-8601, Japan

SOURCE: Parasitology (2007), 134(11), 1639-1647

CODEN: PARAAE; ISSN: 0031-1820

PUBLISHER: Cambridge University Press

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Procyclic forms of *Trypanosoma brucei brucei* remain and propagate in the midgut of tsetse fly where iron is rich. Addnl. iron is also required for their growth in in vitro culture. However, little is known about the genes involved in iron metabolism and the mechanism of iron utilization in procyclic-form cells. Therefore, we surveyed the genes involved in iron metabolism in the T. b. brucei genome sequence database. We found a potential homolog of vacuole protein sorting 41 (VPS41), a gene that is required for high-affinity iron transport in *Saccharomyces cerevisiae* and cloned the full-length gene (TbVPS41). Complementation anal. of TbVPS41 in ΔScvps41 yeast cells showed that TbVPS41 could partially suppress the inability of ΔScvps41 yeast cells to grow on low-iron medium, but it could not suppress the fragmented vacuole phenotype. Further RNA interference (RNAi)-mediated gene knock-down in procyclic-form cells resulted in a significant reduction of growth in low-iron medium; however, no change in growth was observed in normal culture medium. Transmission electron microscopy showed that RNAi caused T. b. brucei cells to have larger nos. of small intracellular vesicles, similar to the fragmented vacuoles observed in ΔScvps41 yeast cells. The present study demonstrates that TbVPS41 plays an important role in the intracellular

iron utilization system as well as in the maintenance of normal cellular morphol.

REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 6 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2007:476913 CAPLUS Full-text

DOCUMENT NUMBER: 147:249690

TITLE: Advances in drug discovery and biochemical studies

AUTHOR(S): Kita, Kiyoshi; Shiomi, Kazuro; Omura, Satoshi

CORPORATE SOURCE: Department of Biomedical Chemistry, Graduate School of Medicine, University of Tokyo, Tokyo, 113-0033, Japan

SOURCE: Trends in Parasitology (2007), 23(5), 223-229

CODEN: TPRACT; ISSN: 1471-4922

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review. Japanese researchers continue to discover new means to combat parasites and make important contributions toward developing tools for global control of parasitic diseases. Streptomyces avermectinius, the source of ivermectin, was discovered in Japan in the early 1970s and renewed and vigorous screening of microbial metabolites in recent years has led to the discovery of new antiprotozoals and anthelmintics, including antimalarial drugs. Intensive studies of parasite energy metabolism, such as NADH-fumarate reductase systems and the synthetic pathways of nucleic acids and amino acids, also contribute to the identification of novel and unique drug targets.

REFERENCE COUNT: 72 THERE ARE 72 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 7 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2006:58924 CAPLUS Full-text

DOCUMENT NUMBER: 145:39855

TITLE: Chemotherapeutic efficacy of ascofuranone in

Trypanosoma vivax-infected mice without glycerol

AUTHOR(S): Yabu, Yoshisada; Suzuki, Takashi; Nihei, Coh-ichi;

Minagawa, Nobuko; Hosokawa, Tomoyoshi; Nagai, Kazuo;

Kita, Kiyoshi; Ohta, Nobuo

CORPORATE SOURCE: Department of Molecular Parasitology, Graduate School of Medical Sciences, Nagoya City University, Nagoya, 467-8601, Japan

SOURCE: Parasitology International (2006), 55(1), 39-43

CODEN: PAINFD; ISSN: 1383-5769

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Ascofuranone, an antibiotic isolated from Ascochyta visiae, showed trypanocidal activity in Trypanosoma vivax-infected mice. A single dose of 50 mg/kg ascofuranone effectively cured the mice without the help of glycerol. Repeated administrations of this drug further enhanced its chemotherapeutic effect. After two, three, and four consecutive days treatment, the doses needed to cure the infection decreased to 25, 12, and 6 mg/kg, so that the total doses administered were 50, 36 and 24 mg/kg, resp. Ascofuranone (50 mg/kg) also had a prophylactic effect against T. vivax infection within the first two days after administration. This prophylactic activity diminished to 80% by day 3 and completely disappeared four days after administration. Of particular interest in this study was that ascofuranone had trypanocidal activity in T. vivax-infected mice in the absence of glycerol, whereas co-administration of glycerol or repeated administrations of this drug are needed for Trypanosoma brucei brucei infection. Our present results strongly suggest

that ascofuranone is also an effective tool in chemotherapy against African trypanosomiasis in domestic animals.

REFERENCE COUNT: 59 THERE ARE 59 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 8 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2006:58922 CAPLUS Full-text

DOCUMENT NUMBER: 145:285774

TITLE: Genetic diversity and kinetic properties of *Trypanosoma cruzi* dihydroorotate dehydrogenase isoforms

AUTHOR(S): Sariego, Idalia; Annoura, Takeshi; Nara, Takeshi; Hashimoto, Muneaki; Tsubouchi, Akiko; Iizumi, Kyoichi; Makiuchi, Takashi; Murata, Eri; Kita, Kiyoshi; Aoki, Takashi

CORPORATE SOURCE: Department of Molecular and Cellular Parasitology, Juntendo University School of Medicine, Hongo 2-1-1, Bunkyo-ku, Tokyo, 113-8421, Japan

SOURCE: Parasitology International (2006), 55(1), 11-16  
CODEN: PAINFJ; ISSN: 1383-5769

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Dihydroorotate dehydrogenase (DHOD) is the fourth enzyme in the de novo pyrimidine biosynthetic pathway and is essential in *Trypanosoma cruzi*, the parasitic protist causing Chagas' disease. *T. cruzi* and human DHOD have different biochem. properties, including the electron acceptor capacities and cellular localization, suggesting that *T. cruzi* DHOD may be a potential chemotherapeutic target against Chagas' disease. Here, we report nucleotide sequence polymorphisms of *T. cruzi* DHOD genes and the kinetic properties of the recombinant enzymes. *T. cruzi* Tulahuen strain possesses three DHOD genes: DHOD1 and DHOD2, involved in the pyrimidine biosynthetic (pyr) gene cluster on an 800 and a 1000 kb chromosomal DNA, resp., and DHOD3, located on an 800 kb DNA. The open reading frames of all three DHOD genes are comprised of 942 bp, and encode proteins of 314 amino acids. The three DHOD genes differ by 26 nucleotides, resulting in replacement of 8 amino acid residues. In contrast, all residues critical for constituting the active site are conserved among the three proteins. Recombinant *T. cruzi* DHOD1 and DHOD2 expressed in *E. coli* possess similar enzymic properties, including optimal pH, optimal temperature, Vmax, and Km for dihydroorotate and fumarate. In contrast, DHOD3 had a higher Vmax and Km for both substrates. Orotate competitively inhibited all three DHOD enzymes to a comparable level. These results suggest that, despite their genetic variations, kinetic properties of the three *T. cruzi* DHODs are conserved. Our findings facilitate further exploitation of *T. cruzi* DHOD inhibitors, as chemotherapeutic agents against Chagas' disease.

REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 9 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:1227142 CAPLUS Full-text

DOCUMENT NUMBER: 144:48333

TITLE: mtDNA of parasites

AUTHOR(S): Watanabe, Yoh-ichi; Kita, Kiyoshi

CORPORATE SOURCE: Grad. Sch. Med., The Univ. Tokyo, Japan

SOURCE: Tanpakushitsu Kakusan Koso (2005), 50(14, Zokan), 1817-1821

CODEN: TAKKAJ; ISSN: 0039-9450

PUBLISHER: Kyoritsu Shuppan

DOCUMENT TYPE: Journal; General Review

LANGUAGE: Japanese

AB A review on mtDNA, codon, functional RNA, mitochondrial protein synthesis, RNA editing, etc., in helminth, trypanosoma, and Plasmodium falciparum.

L45 ANSWER 10 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:1062216 CAPLUS Full-text

DOCUMENT NUMBER: 144:288361

TITLE: Expression, purification and crystallization of Trypanosoma cruzi dihydroorotate dehydrogenase complexed with orotate

AUTHOR(S): Inaoka, Daniel Ken; Takashima, Eizo; Osanai, Arihiro; Shimizu, Hironari; Nara, Takeshi; Aoki, Takashi; Harada, Shigeharu; Kita, Kiyoshi

CORPORATE SOURCE: Department of Biomedical Chemistry, Graduate School of Medicine, University of Tokyo, Bunkyo-ku, Tokyo, 113-0033, Japan

SOURCE: Acta Crystallographica, Section F: Structural Biology and Crystallization Communications (2005), F61(10), 875-878

CODEN: ACSFCL; ISSN: 1744-3091

PUBLISHER: Blackwell Publishing Ltd.

DOCUMENT TYPE: Journal; (online computer file)

LANGUAGE: English

AB Dihydroorotate dehydrogenase (DHOD) catalyzes the oxidation of dihydroorotate to orotate, the 4th step and the only redox reaction in the de novo biosynthesis of pyrimidine. Here, DHOD of T. cruzi (TcDHOD) was expressed as a recombinant protein in Escherichia coli and purified to homogeneity. Crystals of the TcDHOD-orotate complex were grown at 277 K by the sitting-drop vapor-diffusion technique using polyethylene glycol 3350 as precipitant. The crystals diffracted to better than 1.8 Å resolution using synchrotron radiation ( $\lambda = 0.900$  Å). X-ray diffraction data were collected at 100 K and processed to 1.9 Å resolution with 98.2% completeness and an overall Rmerge of 7.8%. The TcDHOD crystals belonged to orthorhombic space group P212121, with unit-cell parameters  $a = 67.87$ ,  $b = 71.89$ , and  $c = 123.27$  Å. The presence of 2 mols. in the asym. unit ( $2 \times 34$  kDa) gave a crystal volume per protein weight (VM) of  $2.2 \text{ Å}^3 \text{ Da}^{-1}$  and a solvent content of 44%.

REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 11 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:647389 CAPLUS Full-text

DOCUMENT NUMBER: 143:244041

TITLE: Mutational analysis of the Trypanosoma vivax alternative oxidase: The E(X)6Y motif is conserved in both mitochondrial alternative oxidase and plastid terminal oxidase and is indispensable for enzyme activity

AUTHOR(S): Nakamura, Kosuke; Sakamoto, Kimitoshi; Kido, Yasutoshi; Fujimoto, Yoko; Suzuki, Takashi; Suzuki, Mitsuko; Yabu, Yoshisada; Ohta, Nobuo; Tsuda, Akiko; Onuma, Misao; Kita, Kiyoshi

CORPORATE SOURCE: Graduate School of Medicine, Department of Biomedical Chemistry, The University of Tokyo, Tokyo, 113-0033, Japan

SOURCE: Biochemical and Biophysical Research Communications (2005), 334(2), 593-600

CODEN: BBRCA9; ISSN: 0006-291X

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Based on amino acid sequence similarity and the ability to catalyze the four-electron reduction of oxygen to water using a quinol substrate, mitochondrial alternative oxidase (AOX) and plastid terminal oxidase (PTOX) appear to be two closely related members of the membrane-bound diiron carboxylate group of proteins. In the current studies, we took advantage of the high activity of *Trypanosoma vivax* AOX (TvAOX) to examine the importance of the conserved Glu and the Tyr residues around the predicted third helix region of AOXs and PTOXs. We first compared the amino acid sequences of TvAOX with AOXs and PTOXs from various taxa and then performed alanine-scanning mutagenesis of TvAOX between amino acids Y199 and Y247. We found that the ubiquinol oxidase activity of TvAOX is completely lost in the E214A mutant, whereas mutants E215A and E216A retained more than 30% of the wild-type activity. Among the Tyr mutants, a complete loss of activity was also observed for the Y221A mutant, whereas the activities were equivalent to wild-type for the Y199A, Y212A, and Y247A mutants. Finally, residues Glu214 and Tyr221 were found to be strictly conserved among AOXs and PTOXs. Based on these findings, it appears that AOXs and PTOXs are a novel subclass of diiron carboxylate proteins that require the conserved motif E(X)6Y for enzyme activity.

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 12 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:993083 CAPLUS Full-text

DOCUMENT NUMBER: 144:287178

TITLE: Alternative oxidase (AOX) genes of African trypanosomes: phylogeny and evolution of AOX and plastid terminal oxidase families

AUTHOR(S): Suzuki, Takashi; Hashimoto, Tetsuo; Yabu, Yoshisada; Majiwa, Phelix A. O.; Ohshima, Shigeru; Suzuki, Mitsuko; Lu, Shaohong; Hato, Mariko; Kido, Yasutoshi; Sakamoto, Kimitoshi; Nakamura, Kosuke; Kita, Kiyoshi; Ohta, Nobuo

CORPORATE SOURCE: Department of Molecular Parasitology, Graduate School of Medical Sciences, Nagoya City University, Kawasumi, Mizuho, Nagoya, 467-8601, Japan

SOURCE: Journal of Eukaryotic Microbiology (2005), 52(4), 374-381

CODEN: JEMIED; ISSN: 1066-5234

PUBLISHER: Blackwell Publishing, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB To clarify evolution and phylogenetic relationships of trypanosome alternative oxidase (AOX) mols., AOX genes (cDNAs) of the African trypanosomes, *Trypanosoma congolense* and *Trypanosoma evansi*, were cloned by PCR. Both AOXs possess conserved consensus motifs (-E-, -EXXH-). The putative amino acid sequence of the AOX of *T. evansi* was exactly the same as that of *T. brucei*. A protein phylogeny of trypanosome AOXs revealed that three genetically and pathogenically distinct strains of *T. congolense* are closely related to each other. When all known AOX sequences collected from current databases were analyzed, the common ancestor of these three *Trypanosoma* species shared a sister-group position to *T. brucei*/*T. evansi*. Monophyly of *Trypanosoma* spp. was clearly supported (100% bootstrap value) with *Trypanosoma vivax* placed at the most basal position of the *Trypanosoma* clade. Monophyly of other eukaryotic lineages, terrestrial plants + red algae, Metazoa, diatoms, Alveolata, oomycetes, green algae, and Fungi, was reconstructed in the best AOX tree obtained from maximum likelihood anal., although some of these clades were not strongly supported. The terrestrial plants + red algae clade showed the closest affinity with an  $\alpha$ -proteobacterium. *Novosphingobium aromaticivorans*, and the common ancestor of these lineages, was separated from

other eukaryotes. Although the root of the AOX subtree was not clearly determined, subsequent phylogenetic anal. of the composite tree for AOX and plastid terminal oxidase (PTOX) demonstrated that PTOX and related cyanobacterial sequences are of a monophyletic origin and their common ancestor is linked to AOX sequences.

REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 13 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2004:8554 CAPLUS Full-text

DOCUMENT NUMBER: 140:232201

TITLE: Direct evidence for cyanide-insensitive quinol oxidase (alternative oxidase) in apicomplexan parasite *Cryptosporidium parvum*: phylogenetic and therapeutic implications

AUTHOR(S): Suzuki, Takashi; Hashimoto, Tetsuo; Yabu, Yoshisada; Kido, Yasutoshi; Sakamoto, Kimitoshi; Nihei, Coh-ichi; Hato, Mariko; Suzuki, Shu-ichi; Amano, Yuko; Nagai, Kazuo; Hosokawa, Tomoyoshi; Minagawa, Nobuko; Ohta, Nobuo; Kita, Kiyoshi

CORPORATE SOURCE: Graduate School of Medical Sciences, Department of Molecular Parasitology, Nagoya City University, Nagoya, 467-8601, Japan

SOURCE: Biochemical and Biophysical Research Communications (2004), 313(4), 1044-1052  
CODEN: BBRCA9; ISSN: 0006-291X

PUBLISHER: Elsevier Science

DOCUMENT TYPE: Journal

LANGUAGE: English

AB *Cryptosporidium parvum* is a parasitic protozoan that causes the diarrheal disease cryptosporidiosis, for which no satisfactory chemotherapy is currently available. Although the presence of mitochondria in this parasite has been suggested, its respiratory system is poorly understood due to difficulties in performing biochem. analyses. In order to better understand the respiratory chain of *C. parvum*, we surveyed its genomic DNA database in GenBank and identified a partial sequence encoding cyanide-insensitive alternative oxidase (AOX). Based on this sequence, we cloned *C. parvum* AOX (CpAOX) cDNA from the phylum Apicomplexa for the first time. The deduced amino acid sequence (335 a.a.) of CpAOX contains diiron coordination motifs (-E-, -EXXH-) that are conserved among AOXs. Phylogenetic anal. suggested that CpAOX is a mitochondrial-type AOX, possibly derived from mitochondrial endosymbiont gene transfer. The recombinant enzyme expressed in *Escherichia coli* showed quinol oxidase activity. This activity was insensitive to cyanide and highly sensitive to ascofuranone, a specific inhibitor of trypanosome AOX.

REFERENCE COUNT: 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 14 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2004:885782 CAPLUS Full-text

DOCUMENT NUMBER: 141:374297

TITLE: Drug development at global level

AUTHOR(S): Kita, Kiyoshi

CORPORATE SOURCE: Grad. Sch. Med., The Univ. Tokyo, Japan

SOURCE: Farumashia (2004), 40(10), 909-913  
CODEN: FARUAW; ISSN: 0014-8601

PUBLISHER: Pharmaceutical Society of Japan

DOCUMENT TYPE: Journal; General Review

LANGUAGE: Japanese

AB A review on the development of parasiticides, especially, ascofuranone, which inhibits trypanosome alternative oxidase for control of *Trypanosoma brucei* in treatment of African trypanosomiasis.

L45 ANSWER 15 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2004:404924 CAPLUS Full-text

DOCUMENT NUMBER: 141:200963

TITLE: Molecular cloning and characterization of *Trypanosoma vivax* alternative oxidase (AOX) gene, a target of the trypanocide ascofuranone

AUTHOR(S): Suzuki, Takashi; Nihei, Coh-Ichi; Yabu, Yoshisada; Hashimoto, Tetsuo; Suzuki, Mitsuko; Yoshida, Ayako; Nagai, Kazuo; Hosokawa, Tomoyoshi; Minagawa, Nobuko; Suzuki, Shuichi; Kita, Kiyoshi; Ohta, Nobuo

CORPORATE SOURCE: Graduate School of Medical Sciences, Department of Molecular Parasitology, Nagoya City University, Nagoya, 467-8601, Japan

SOURCE: Parasitology International (2004), 53(3), 235-245  
CODEN: PAINFD; ISSN: 1383-5769

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

AB *Trypanosoma vivax* causes nagana disease in cattle. Since *T. vivax* is transmitted not only by tsetse flies but also by other biting flies (non-cyclic transmission), the parasite has been distributed to and has had a significant economic impact on wide geog. areas, including Africa and South America. Our previous study on *Trypanosoma brucei brucei* showed that the trypanosome alternative oxidase (TAO, TbAOX) is a promising target of chemotherapy. For this reason, we also have cloned the *T. vivax* AOX (TvAOX) gene and characterized the recombinant enzyme. The deduced amino acid sequence (328 a.a.) of TvAOX shares 76% identity with TbAOX and contains the diiron-coordination motifs (-E-, -EXXH-) that are conserved among AOXs. The  $K_m$  of recombinant TvAOX (rTvAOX) expressed in *Escherichia coli* for ubiquinol ( $87.0 \pm 0.54 \mu M$ ) was significantly lower than the value for recombinant TbAOX (rTbAOX) ( $714 \pm 4.5 \mu M$ ). Ascofuranone, the most potent inhibitor of TbAOX, was a competitive inhibitor of rTvAOX with a  $K_i$  value ( $0.40 \pm 0.00$  nM) significantly lower than that for rTbAOX ( $1.29 \pm 0.00$  nM). The non-cyclic transmission ability of *T. vivax* and the in vivo chemotherapeutic efficacy of ascofuranone against *T. vivax* and *T. b. brucei* infection are discussed in terms of these  $K_m$  and  $K_i$  values.

REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 16 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:926536 CAPLUS Full-text

DOCUMENT NUMBER: 140:209735

TITLE: Parasite mitochondria as drug target: Diversity and dynamic changes during the life cycle

AUTHOR(S): Kita, Kiyoshi; Nihei, Coichi; Tomitsuka, Eriko

CORPORATE SOURCE: Department of Biomedical Chemistry, Graduate School of Medicine, University of Tokyo, Tokyo, 113-0033, Japan

SOURCE: Current Medicinal Chemistry (2003), 10(23), 2535-2548  
CODEN: CMCHE7; ISSN: 0929-8673

PUBLISHER: Bentham Science Publishers Ltd.

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review. Parasites have developed a wide variety of physiol. functions to survive within the specialized environments of the host. Regarding energy

metabolism, which represents an essential factor for survival, parasites adapt low oxygen tension in host mammals using metabolic systems that differ substantially from those of the host. Most parasites do not use free oxygen available within the host, but employ systems other than oxidative phosphorylation for ATP synthesis. Furthermore, parasites display marked changes in mitochondrial morphol. and components during the life cycle, and these represent very interesting elements of biol. processes such as developmental control and environmental adaptation. The enzymes in parasite-specific pathways offer potential targets for chemotherapy. Cyanide-insensitive trypanosome alternative oxidase (TAO) is the terminal oxidase of the respiratory chain of long slender bloodstream forms of the African trypanosome, which causes sleeping sickness. Recently, the most potent inhibitor of TAO to date, ascofuranone, was isolated from the phytopathogenic fungus, *Ascochyta visiae*. The inhibitory mechanisms of ascofuranone have been revealed using recombinant enzyme. Parasite-specific respiratory systems are also found in helminths. The NADH-fumarate reductase system in mitochondria form a final step in the phosphoenolpyruvate carboxykinase (PEPCK)-succinate pathway, which plays an important role in anaerobic energy metabolism for the *Ascaris suum* adult. Enzymes in this system, such as NADH-rhodoquinone reductase (complex I) and rhodoquinol-fumarate reductase (complex II), form promising targets for chemotherapy. In fact, a specific inhibitor of nematode complex I, nafuredin, has been found in mass-screening using parasite mitochondria.

REFERENCE COUNT: 107 THERE ARE 107 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 17 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:822795 CAPLUS Full-text

DOCUMENT NUMBER: 139:357611

TITLE: Ascofuranone as a chemotherapeutic agent of African sleeping sickness

AUTHOR(S): Yabu, Yoshisada; Suzuki, Takashi; Kita, Kiyoshi

CORPORATE SOURCE: Dep. Mol. Parasitol., Nagoya City Univ., Nagoya, 467-8601, Japan

SOURCE: Baio Saiensu to Indasutori (2003), 61(10), 681-682  
CODEN: BIDSE6; ISSN: 0914-8981

PUBLISHER: Baioindasutori Kyokai

DOCUMENT TYPE: Journal; General Review

LANGUAGE: Japanese

AB A review on African sleeping sickness caused by *Trypanosoma* infection, inhibition of trypanosome alternative oxidase by ascofuranone, and therapeutic effect of ascofuranone in African sleeping sickness mouse models.

L45 ANSWER 18 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:621706 CAPLUS Full-text

DOCUMENT NUMBER: 139:335515

TITLE: Metabolic characteristics in parasites

AUTHOR(S): Kita, Kiyoshi

CORPORATE SOURCE: Graduate School of Medicine, University of Tokyo, Japan

SOURCE: Chiryogaku (2003), 37(6), 592-596  
CODEN: CHRYDT; ISSN: 0386-8109

PUBLISHER: Raifu Saiensu Shuppan K.K.

DOCUMENT TYPE: Journal; General Review

LANGUAGE: Japanese

AB A review, on energy metabolism and related enzymes in parasites, discussing the metabolism of nucleic acids, amino acids, and lipids in parasites, such as *Ascaris suum*, and *Trypanosoma cruzi*.



L45 ANSWER 19 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:787439 CAPLUS Full-text

DOCUMENT NUMBER: 140:180196

TITLE: Overproduction of highly active trypanosome alternative oxidase in Escherichia coli heme-deficient mutant

AUTHOR(S): Fukai, Yoshihisa; Nihei, Coichi; Kawai, Keisuke; Yabu, Yoshisada; Suzuki, Takasi; Ohta, Nobuo; Minagawa, Nobuko; Nagai, Kazuo; Kita, Kiyoshi

CORPORATE SOURCE: Department of Biomedical Chemistry, Graduate School of Medicine, University of Tokyo, Bunkyo-ku, Tokyo, 113-0033, Japan

SOURCE: Parasitology International (2003), 52(3), 237-241  
CODEN: PAINFD; ISSN: 1383-5769

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Cyanide-insensitive trypanosome alternative oxidase (TAO) is the terminal oxidase of the respiratory chain of long slender bloodstream forms of the African trypanosome, which causes sleeping sickness in humans and nagana in cattle. TAO has been targeted for the development of anti-trypanosomal drugs, because it does not exist in the host. In this study, we established a system for overprodn. of highly active TAO in Escherichia coli heme-deficient mutant. Kinetic anal. of recombinant enzyme and TAO in Trypanosoma brucei brucei mitochondria revealed that recombinant TAO retains the properties of native enzyme, indicating that recombinant TAO is quite valuable for further biochem. study of TAO.

REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 20 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:443379 CAPLUS Full-text

DOCUMENT NUMBER: 140:22662

TITLE: The efficacy of ascofuranone in a consecutive treatment on Trypanosoma brucei brucei in mice

AUTHOR(S): Yabu, Yoshisada; Yoshida, Ayako; Suzuki, Takashi; Nihei, Coh-ichi; Kawai, Keisuke; Minagawa, Nobuko; Hosokawa, Tomoyoshi; Nagai, Kazuo; Kita, Kiyoshi; Ohta, Nobuo

CORPORATE SOURCE: Department of Molecular Parasitology, Nagoya City University, Nagoya, 467-8601, Japan

SOURCE: Parasitology International (2003), 52(2), 155-164  
CODEN: PAINFD; ISSN: 1383-5769

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Consecutive administration of ascofuranone without glycerol was found to have therapeutic efficacy against Trypanosoma brucei brucei infection in mice. A suspension of ascofuranone (25-100 mg/kg) was administered i.p. every 24 h for 1-4 consecutive days to trypanosome-infected mice and efficacy was compared with oral treatment. With i.p. administration, all mice treated with 100 mg/kg ascofuranone for 4 consecutive days were cured. On contrary, with oral treatment a higher dose of ascofuranone (400 mg/kg) was needed for 8 consecutive days to cure the mice. With i.p. treatment, parasitemia was strongly suppressed, with almost all long slender bloodstream forms of the parasite changed to short stumpy forms by day 3 and the parasites were eliminated 4 days after the start of treatment. These ascofuranone-induced short stumpy forms were morphol. analogous to the stumpy forms 2 days after

peak parasitemia of pleomorphic clone of *T. b. brucei* GUTat 3.1. However, the properties of ubiquinol oxidase activity, which is the target of ascofuranone, in mitochondria isolated from before and after treatment, were almost same. The enzymic activities of ubiquinol oxidase were only decreased to approx. 30% within a day after treatment, and then kept at nearly the same level. In the present study, we have improved the regimen for administration of ascofuranone without glycerol, and demonstrated that consecutively administered ascofuranone showed trypanocidal effects in *T. b. brucei* infected mice. Our present results strongly suggest that consecutive administration of ascofuranone may be an effective chemotherapy for African trypanosomiasis.

REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 21 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:196106 CAPLUS Full-text

DOCUMENT NUMBER: 139:32393

TITLE: Purification of active recombinant trypanosome alternative oxidase

AUTHOR(S): Nihei, Coichi; Fukai, Yoshihisa; Kawai, Keisuke; Osanai, Arihiro; Yabu, Yoshisada; Suzuki, Takashi; Ohta, Nobuo; Minagawa, Nobuko; Nagai, Kazuo; Kita, Kiyoshi

CORPORATE SOURCE: Graduate School of Medicine, Department of Biomedical Chemistry, University of Tokyo, Bunkyo-ku, Tokyo, 113-0033, Japan

SOURCE: FEBS Letters (2003), 538(1-3), 35-40  
CODEN: FEBLAL; ISSN: 0014-5793

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Trypanosome alternative oxidase (TAO) is the terminal oxidase of the respiratory chain in long slender bloodstream forms of African trypanosomes. TAO is a cytochrome-independent, cyanide-insensitive quinol oxidase. These characteristics are distinct from those of the bacterial quinol oxidases, proteins that belong to the heme-copper terminal oxidase superfamily. The inability to purify stable TAO has severely hampered biochem. studies of the alternative oxidase family. In the present study, we were able to purify recombinant TAO to homogeneity from *Escherichia coli* membranes using the detergent digitonin. Kinetic anal. of the purified TAO revealed that the specific inhibitor ascofuranone is a competitive inhibitor of ubiquinol oxidase activity.

REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 22 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2002:455276 CAPLUS Full-text

DOCUMENT NUMBER: 138:49228

TITLE: Target molecule of the novel anti-Trypanosoma drug ascofuranone: trypanosome alternative oxidase

AUTHOR(S): Nihei, Koichi; Kita, Kiyoshi

CORPORATE SOURCE: Graduate School of Medicine, University of Tokyo, Japan

SOURCE: Kagaku to Kyoiku (2002), 50(5), 350-354  
CODEN: KAKYKY; ISSN: 0386-2151

PUBLISHER: Nippon Kagakkai

DOCUMENT TYPE: Journal; General Review

LANGUAGE: Japanese

AB A review, discussing the action mechanism and pharmacol. of the anti-Trypanosoma drug ascofuranone against *Trypanosoma brucei* infestation by targeting trypanosome alternative oxidase.

L45 ANSWER 23 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:279181 CAPLUS Full-text

DOCUMENT NUMBER: 139:97132

TITLE: Purification of recombinant trypanosome alternative oxidase

AUTHOR(S): Kawai, K.; Nihei, C.; Fukai, Y.; Yabu, Y.; Ohta, N.; Minagawa, N.; Nagai, K.; Kita, K.

CORPORATE SOURCE: Department of Biomedical Chemistry, Graduate School of Medicine, The University of Tokyo, Japan

SOURCE: Parasitology--ICOPA X: Symposia, Workshops and Contributed Papers, Proceedings of the International Congress, 10th, Vancouver, BC, Canada, Aug. 4-9, 2002 (2002), 295-301. Monduzzi Editore: Bologna, Italy. CODEN: 69DTB8; ISBN: 88-323-2804-6

DOCUMENT TYPE: Conference

LANGUAGE: English

AB Trypanosome alternative oxidase (TAO) is the terminal oxidase of the respiratory chain of long slender bloodstream forms (LS forms) of African trypanosome, which causes sleeping sickness in human and nagana in cattle. TAO is a cytochrome-independent, cyanide-insensitive quinol oxidase and these properties are quite different from those of the bacterial quinol oxidase which belongs to the heme-copper terminal oxidase superfamily. Only little information concerning the mol. structure and enzymic features of TAO have been available, whereas the bacterial enzyme has been well characterized. In the present study, we constructed an E. coli expression system for TAO. The recombinant TAO (rTAO) was expressed in E. coli  $\Delta$ hemA mutant, FN 102/pTAO and purified from the membrane of the E. coli to homogeneity.

REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 24 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2002:476087 CAPLUS Full-text

DOCUMENT NUMBER: 138:82708

TITLE: Trypanosome alternative oxidase as a target of chemotherapy

AUTHOR(S): Nihei, Coichi; Fukai, Yoshihisa; Kita, Kiyoshi

CORPORATE SOURCE: Graduate School of Medicine, Department of Biomedical Chemistry, University of Tokyo, Bunkyo-ku, Tokyo, 113-0033, Japan

SOURCE: Biochimica et Biophysica Acta, Molecular Basis of Disease(2002), 1587(2-3), 234-239  
CODEN: BBADEX; ISSN: 0925-4439

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review. Parasites have developed a variety of physiol. functions necessary for their survival within the specialized environment of the host. Using metabolic systems that are very different from those of the host, they can adapt to low oxygen tension present within the host animals. Most parasites do not use the oxygen available within the host to generate ATP, but rather employ systems anaerobic metabolic pathways. The enzymes in these parasite-specific pathways are potential targets for chemotherapy. Cyanide-insensitive trypanosome alternative oxidase (TAO) is the terminal oxidase of the respiratory chain of long slender bloodstream forms of the African trypanosome, which causes sleeping sickness in human and nagana in cattle. TAO has been targeted for the development of anti-trypanosomal drugs because it does not exist in the host. Recently, we found the most potent inhibitor

of TAO to date, ascofuranone, a compound isolated from the phytopathogenic fungus, *Ascochyta visiae*.

REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 25 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2002:511240 CAPLUS Full-text

DOCUMENT NUMBER: 138:1625

TITLE: Strain-specific difference in amino acid sequences of trypanosome alternative oxidase

AUTHOR(S): Fukai, Yoshihisa; Nihei, Coichi; Yabu, Yoshisada; Suzuki, Takasi; Ohta, Nobuo; Minagawa, Nobuko; Nagai, Kazuo; Kita, Kiyoshi

CORPORATE SOURCE: Graduate School of Medicine, Department of Biomedical Chemistry, The University of Tokyo, Bunkyo-ku, Tokyo, 113-0033, Japan

SOURCE: Parasitology International (2002), 51(2), 195-199  
CODEN: PAINFD; ISSN: 1383-5769

PUBLISHER: Elsevier Science Ireland Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Cyanide-insensitive trypanosome alternative oxidase (TAO) is the terminal oxidase of the respiratory chain of long slender bloodstream forms of the African trypanosome, which causes sleeping sickness in human and nagana in cattle. TAO has been targeted for the development of anti-trypanosomal drugs because it does not exist in the host. The cDNA for TAO has been cloned from *Trypanosoma brucei brucei* EATRO110 strain and has been used for further characterization. In this study, we found amino acid sequence of the C-terminal part of TAO from the strain that we are using, *T. b. brucei* TC221, is considerably different from that of the EATRO110 strain.

REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 26 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2002:509982 CAPLUS Full-text

DOCUMENT NUMBER: 137:290812

TITLE: Characterization of the dihydroorotate dehydrogenase as a soluble fumarate reductase in *Trypanosoma cruzi*

AUTHOR(S): Takashima, Eizo; Inaoka, Daniel Ken; Osanai, Arihiro; Nara, Takeshi; Odaka, Masao; Aoki, Takashi; Inaka, Kozi; Harada, Shigeharu; Kita, Kiyoshi

CORPORATE SOURCE: Department of Biomedical Chemistry, The University of Tokyo, Graduate School of Medicine, Bunkyo-ku, Tokyo, 113-0033, Japan

SOURCE: Molecular and Biochemical Parasitology (2002), 122(2), 189-200

CODEN: MBIPDP; ISSN: 0166-6851

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB *Trypanosoma cruzi*, a protozoan causing Chagas' disease, excretes a considerable amount of succinate even though it uses the TCA cycle and the aerobic respiratory chain. For this reason, it was believed that unknown metabolic pathways participate in succinate production in this parasite. In the present study, we examined the mol. properties of dihydroorotate dehydrogenase (DHOD), the fourth enzyme of de novo pyrimidine biosynthetic pathway, as a soluble fumarate reductase (FRD) because our sequence anal. of pyr genes cluster showed that the amino acid sequence of *T. cruzi* DHOD is quite similar to that of type 1A DHOD of *Saccharomyces cerevisiae*, an enzyme that uses fumarate as an electron acceptor and produces succinate. Biochem.

analyses of the cytosolic enzyme purified from the parasite and of the recombinant enzyme revealed that *T. cruzi* DHOD has methylviologen-fumarate reductase (MV-FRD) activity. In addition, *T. cruzi* DHOD was found to catalyze electron transfer from dihydroorotate to fumarate by a ping-pong Bi-Bi mechanism. The recombinant enzyme contained FMN as a prosthetic group. Dynamic light scattering anal. indicated that *T. cruzi* DHOD is a homodimer. These results clearly indicated that the cytosolic MV-FRD is attributable to *T. cruzi* DHOD. The DHOD may play an important role in succinate/fumarate metabolism as well as de novo pyrimidine biosynthesis in *T. cruzi*.

REFERENCE COUNT: 48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 27 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2002:103671 CAPLUS Full-text

DOCUMENT NUMBER: 136:98883

TITLE: Adaptation to low oxygen tension in parasite mitochondria

AUTHOR(S): Kita, Kiyoshi

CORPORATE SOURCE: Grad. Sch. Med., The Univ. Tokyo, Japan

SOURCE: Tanpakushitsu Kakusan Koso (2002), 47(1), 37-44

CODEN: TAKKAJ; ISSN: 0039-9450

PUBLISHER: Kyoritsu Shuppan

DOCUMENT TYPE: Journal; General Review

LANGUAGE: Japanese

AB A review on the functions of mitochondria of protozoa, changes of mitochondrial functions of *Trypanosoma brucei brucei* during life cycle, structure and function of trypanosome alternative oxidase (cyanide-insensitive quinol oxidase), inhibition of glycerol-3-phosphate-dependent mitochondrial O<sub>2</sub> consumption by antitrypanosome drug ascofuranone, changes of energy metabolism and respiratory chain in *Ascaris suum* in response to oxygen tension during life-cycle, NADH-fumarate reductase system of adult *A. suum*, functions of complex II as a succinate-ubiquinone reductase in larvae and as a quinol-fumarate reductase in adults, association of mitochondrial quinol-fumarate reductase activity with parasitic adaptation, evolution of quinones, and importance of the parasite mitochondrias as targets of chemotherapeutic agents against parasites.

L45 ANSWER 28 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2001:350991 CAPLUS Full-text

DOCUMENT NUMBER: 135:163655

TITLE: environmental adaptation of the respiratory system in parasites

AUTHOR(S): Kita, Kiyoshi

CORPORATE SOURCE: Graduate School of Medical Research, University of Tokyo, Japan

SOURCE: Shirizu Baiosaiensu no Shinseiki (2000), Volume 7, 47-59. Editor(s): Yoshida, Masasuke; Mogi, Tatsushi.

Kyoritsu Shuppan: Tokyo, Japan.

CODEN: 69BHTE

DOCUMENT TYPE: Conference; General Review

LANGUAGE: Japanese

AB A review with 20 refs., on changes of mitochondrial respiratory chains in parasites (such as *Ascaris suum* and *Trypanosoma brucei brucei*) and their relations to the parasitic life cycle and environmental adaptation.

L45 ANSWER 29 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1999:686146 CAPLUS Full-text

DOCUMENT NUMBER: 132:60821

TITLE: Functional expression of the ascofuranone-sensitive Trypanosoma brucei brucei alternative oxidase in the cytoplasmic membrane of Escherichia coli

AUTHOR(S): Fukai, Y.; Amino, H.; Hirawake, H.; Yabu, Y.; Ohta, N.; Minagawa, N.; Sakajo, S.; Yoshimoto, A.; Nagai, K.; Takamiya, S.; Kojima, S.; Kita, K.

CORPORATE SOURCE: Bunkyo-ku, 7-3-1 Hongo, Graduate School of Medicine, Department of Biomedical Chemistry, The University of Tokyo, Tokyo, Japan

SOURCE: Comparative Biochemistry and Physiology, Part C: Pharmacology, Toxicology & Endocrinology (1999), 124C(2), 141-148  
CODEN: CBPCEE; ISSN: 0742-8413

PUBLISHER: Elsevier Science Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Trypanosome alternative oxidase (TAO) is the terminal oxidase of the respiratory chain of long slender bloodstream forms (LS forms) of African trypanosome, which causes sleeping sickness in human and nagana in cattle. TAO is a cytochrome-independent, cyanide-insensitive quinol oxidase and these properties are quite different from those of the bacterial quinol oxidase which belongs to the heme-copper terminal oxidase superfamily. Only little information concerning the mol. structure and enzymic features of TAO have been available, whereas the bacterial enzyme has been well characterized. In this study, a cDNA encoding TAO from Trypanosoma brucei brucei was cloned into the expression vector pET15b (pTAO) and recombinant TAO was expressed in Escherichia coli. The growth of the transformant carrying pTAO was cyanide-resistant. A peptide with a mol. mass of 37 kDa was found in the cytoplasmic membrane of E. coli, and was recognized by antibodies against plant-type alternative oxidases from Sauromatum guttatum and Hansenula anomala. Both the ubiquinol oxidase and succinate oxidase activities found in the membrane of the transformant were insensitive to cyanide, while those of the control strain, which contained vector alone, were inhibited. This cyanide-insensitive growth of the E. coli carrying pTAO was inhibited by the addition of ascofuranone, a potent and specific inhibitor of TAO ubiquinol oxidase. The ubiquinol oxidase activity of the membrane from the transformant was sensitive to ascofuranone. These results clearly show the functional expression of TAO in E. coli and indicate that ubiquinol-8 in the E. coli membrane is able to serve as an electron donor to the recombinant enzyme and confer cyanide-resistant and ascofuranone-sensitive growth to E. coli. This system will facilitate the biochem. characterization of the novel terminal oxidase, TAO, and the understanding on the mechanism of the trypanocidal effect of ascofuranone.

REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 30 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1998:400327 CAPLUS Full-text

DOCUMENT NUMBER: 129:117469

TITLE: Trypanocidal effects of curcumin in vitro

AUTHOR(S): Nose, Mitsuhiko; Koide, Tatsuo; Ogiwara, Yukio; Yabu, Yoshisada; Ohta, Nobuo

CORPORATE SOURCE: Department of Pharmacognosy and Plant Chemistry, Faculty of Pharmaceutical Sciences, Nagoya City University, Nagoya, 467, Japan

SOURCE: Biological & Pharmaceutical Bulletin (1998), 21(6), 643-645

CODEN: BPBLEO; ISSN: 0918-6158

PUBLISHER: Pharmaceutical Society of Japan

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Searching for antiparasitic agents from natural sources revealed that curcumin is cytotoxic against African trypanosomes in vitro. The LD50 values of curcumin were  $4.77 \pm 0.91 \mu\text{M}$  for blood stream forms and  $46.52 \pm 4.94 \mu\text{M}$  for procyclic forms of *Trypanosoma brucei brucei* (GUTat 3.1 clone).

REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 31 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1998:400313 CAPLUS Full-text

DOCUMENT NUMBER: 129:132403

TITLE: Formation of reactive oxygen intermediates might be involved in the trypanocidal activity of gallic acid

AUTHOR(S): Nose, Mitsuhiko; Koide, Tatsuo; Morikawa, Kyoko; Inoue, Makoto; Ogihara, Yukio; Yabu, Yoshisada; Ohta, Nobuo

CORPORATE SOURCE: Department of Pharmacognosy and Plant Chemistry, Faculty of Pharmaceutical Sciences, Nagoya City University, Nagoya, 467, Japan

SOURCE: Biological & Pharmaceutical Bulletin (1998), 21(6), 583-587

CODEN: BPBLEO; ISSN: 0918-6158

PUBLISHER: Pharmaceutical Society of Japan

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The authors investigated the mechanism of the trypanocidal activity of gallic acid (GA). GA-induced trypanocidal activity was significantly reduced by pretreatment with superoxide dismutase (SOD) and/or catalase. The ESR technique with 5,5-dimethyl-1-pyrroline N-oxide (DMPO) as a spin trapping agent revealed that a DMPO-OH adduct was detected in culture medium containing GA. The intensity of ESR signals of the DMPO-OH adduct was increased in a time dependent manner. SOD also inhibited the formation of GA-induced DMPO-OH adducts. Furthermore, GA enhanced DNA single-strand breaks induced by Fenton reagent. These results suggest the possibility that GA acts as a pro-oxidant for trypanocidal activity.

REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 32 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1999:26547 CAPLUS Full-text

DOCUMENT NUMBER: 130:265982

TITLE: Parasite infection and apoptosis

AUTHOR(S): Kita, Kiyoshi; Shimada, Junko

CORPORATE SOURCE: Graduate School of Medicine, The University of Tokyo, Japan

SOURCE: Igaku no Ayumi (1998), 187(5), 436-440

CODEN: IGAYAY; ISSN: 0039-2359

PUBLISHER: Ishiyaku Shuppan

DOCUMENT TYPE: Journal; General Review

LANGUAGE: Japanese

AB A review with 19 refs., on apoptosis of parasite-infected host cells; infection by *Leishmania*, *Toxoplasma*, and *Trypanosoma*; malaria or *Trypanosoma* infection-induced apoptosis of host immunocytes, and apoptosis of parasite.

L45 ANSWER 33 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1998:522580 CAPLUS Full-text

DOCUMENT NUMBER: 129:270006

TITLE: Oral and intraperitoneal treatment of *Trypanosoma brucei brucei* with a combination of ascofuranone and

glycerol in mice  
AUTHOR(S): Yabu, Yoshisada; Minagawa, Nobuko; Kita, Kiyoshi;  
Nagai, Kazuo; Honma, Masakatsu; Sakajo, Shigeru;  
Koide, Tatsuo; Ohta, Nobuo; Yoshimoto, Akio  
CORPORATE SOURCE: Department of Medical Zoology, Medical School, Nagoya  
City University, Nagoya, 467-8601, Japan  
SOURCE: Parasitology International (1998), 47(2), 131-137  
CODEN: PAINFD; ISSN: 1383-5769  
PUBLISHER: Elsevier Science Ireland Ltd.  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB A suspension of ascofuranone (6-200 mg/kg) was given orally or i.p., and then glycerol (1 g/kg) was administered orally or i.p. at 30-min intervals to mice heavily parasitemic with *T. brucei brucei*. Both orally (100 mg/kg) and i.p. (25 mg/kg) administered ascofuranone, combined with a total dose of 3 g glycerol/kg, produced potent antitrypanosomal activity in infested mice. The trypanocidal activity of ascofuranone was very powerful, and all trypanosomes had disappeared within 30 and 180 min after final i.p. and oral treatment, resp. This combination treatment showed high efficacy and low toxicity. Ascofuranone in combination with glycerol may be an effective tool in chemotherapy for African trypanosomiasis.

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L45 ANSWER 34 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1998:98846 CAPLUS Full-text  
DOCUMENT NUMBER: 128:149258  
TITLE: Trypanocidal effects of gallic acid and related compounds

AUTHOR(S): Koide, Tatsuo; Nose, Mitsuhiko; Inoue, Makoto;  
Ogihara, Yukio; Yabu, Yoshisada; Ohta, Nobuo  
CORPORATE SOURCE: Dep. Pharmacognosy Plant Chemistry, Fac.  
Pharmaceutical Sciences, Nagoya City Univ., Nagoya,  
467, Japan  
SOURCE: Planta Medica (1998), 64(1), 27-30  
CODEN: PLMEAA; ISSN: 0032-0943  
PUBLISHER: Georg Thieme Verlag  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Gallic acid (3,4,5-trihydroxybenzoic acid) is a naturally abundant plant phenolic compound and it is well known as a component of hydrolyzable tannins. We report here that gallic acid and related compds. have trypanocidal activity against *Trypanosoma brucei brucei* (GUTat 3.1) in both the long slender bloodstream forms and the procyclic forms, in vitro. LD50 values of gallic acid are  $46.96 \pm 1.28 \mu\text{M}$  for bloodstream forms and  $30.02 \pm 3.49$  for procyclic forms, resp. A study of structurally related compds. suggested that the pyrogallol moiety could be responsible for this activity.

L45 ANSWER 35 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1997:179701 CAPLUS Full-text  
DOCUMENT NUMBER: 126:180891  
TITLE: An antibiotic, ascofuranone, specifically inhibits respiration and in vitro growth of long slender bloodstream forms of *trypanosoma brucei brucei*.  
[Erratum to document cited in CA125:265065]  
AUTHOR(S): Minagawa, Nobuko; Yabu, Yoshisada; Kita, Kiyoshi;  
Nagai, Kazuo; Ohta, Nobuo; Meguro, Keiichi; Sakajo, Shigeru; Yoshimoto, Akio  
CORPORATE SOURCE: Department of Biochemistry, Niigata College of



Pharmacy, Niigata, 950-21, Japan  
 SOURCE: Molecular and Biochemical Parasitology (1997), 84(2),  
 271-280  
 CODEN: MBIPDP; ISSN: 0166-6851  
 PUBLISHER: Elsevier  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB The errors were not reflected in the abstract or the index entries.

L45 ANSWER 36 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1996:583098 CAPLUS Full-text

DOCUMENT NUMBER: 125:265065

TITLE: An antibiotic, ascofuranone, specifically inhibits  
 respiration and in vitro growth of long slender  
 bloodstream forms of *Trypanosoma brucei brucei*

AUTHOR(S): Minagawa, Nobuko; Yabu, Yoshisada; Kita, Kiyoshi;  
 Nagai, Kazuo; Ohta, Nobuo; Meguro, Keiichi; Sakajo,  
 Shigeru; Yoshimoto, Akio

CORPORATE SOURCE: Department of Biochemistry, Niigata College of  
 Pharmacy, 5-13-2 Kamishin'ei-cho, Niigata, 950-21,  
 Japan

SOURCE: Molecular and Biochemical Parasitology (1996), 81(2),  
 127-136

CODEN: MBIPDP; ISSN: 0166-6851

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Ascofuranone, a prenylphenol antibiotic isolated from a phytopathogenic  
 fungus, *Ascochyta visiae*, strongly inhibited both glucose-dependent cellular  
 respiration and glycerol-3-phosphate-dependent mitochondrial O<sub>2</sub> consumption of  
 long slender bloodstream forms of *Trypanosoma brucei brucei*. This inhibition  
 was suggested to be due to inhibition of the mitochondrial electron-transport  
 system, composed of glycerol-3-phosphate dehydrogenase (EC 1.1.99.5) and  
 plant-like alternative oxidase. Ascofuranone noncompetitively inhibited the  
 reduced coenzyme Q1-dependent O<sub>2</sub> uptake of the mitochondria with respect to  
 ubiquinol (K<sub>i</sub> = 2.38 nM). Therefore, the susceptible site is deduced to be the  
 ubiquinone redox machinery which links the two enzyme activities. Further,  
 ascofuranone in combination with glycerol completely blocked energy  
 production, and potentially inhibited the in vitro growth of the parasite. Our  
 findings suggest that ascofuranone might be a promising candidate for the  
 chemotherapeutic agents of African trypanosomiasis.

L45 ANSWER 37 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1993:78987 CAPLUS Full-text

DOCUMENT NUMBER: 118:78987

TITLE: Inhibition of IgM antibody-mediated aggregation of  
*Trypanosoma gambiense* in the presence of complement

AUTHOR(S): Takayanagi, T.; Kawaguchi, H.; Yabu, Y.; Itoh, M.;  
 Yano, K.

CORPORATE SOURCE: Med. Sch., Nagoya City Univ., Nagoya, 467, Japan

SOURCE: Experientia (1992), 48(10), 1002-6

CODEN: EXPEAM; ISSN: 0014-4754

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The immune reaction was studied between *T. gambiense* and monoclonal IgM mouse  
 antibody at equivalence with or without rabbit complement. Antibody-mediated  
 trypanosome clumps formed in the absence of complement, and were readily  
 dissociated by complement to become free. In the presence of complement, on

the other hand, *T. gambiense* was not aggregated by the antibody. Free parasites adhered readily to cultured peritoneal macrophages. Complement-mediated dissociation of the clumped trypanosomes in the equivalence area released a large number of previously bound surface antigens. These antigens were capable of binding again to fresh IgM antibody. The complement system caused a functional alternation, changing the multivalent nature of the IgM antibody in the immune complex into a univalent one. This phenomenon is of great advantage to the infected host in clearing pathogens in vivo, as it allows more antibodies to attach to trypanosomes and subsequently initiate complement activity.

L45 ANSWER 38 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1987:513991 CAPLUS Full-text

DOCUMENT NUMBER: 107:113991

TITLE: Contribution of the complement system to antibody-mediated binding to *Trypanosoma gambiense* to macrophages

AUTHOR(S): Takayanagi, Tan; Kawaguchi, Hitoshi; Yabu, Yoshisada; Ito, Makoto; Appawu, Maxwell Alex

CORPORATE SOURCE: Med. Sch., Nagoya City Univ., Nagoya, 467, Japan

SOURCE: Journal of Parasitology (1987), 73(2), 333-41  
CODEN: JOPAA2; ISSN: 0022-3395

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The role of complement in the process of binding of trypanosomes to macrophages in the presence of specific antibody was studied. The aggregation of trypanosomes observed at the optimal antigen-antibody ratio or in the presence of excess antigen inhibited the binding. Complement caused clumped trypanosomes to dissociate, and the free trypanosomes, which were presumed to be coated with antibody that had fixed complement, readily attached to surfaces of phagocytes. Thus, complement contributed at the site of the antigen-antibody reaction to the creation of an environment suitable for the binding. Apparently, the trypanosomes dissociated by complement adhered to C3 receptors of the macrophage. However, in the absence of complement and in regions of antibody excess, free trypanosomes also attached to phagocytes. Thus, phagocytes may also have receptors for the Fc portion of aggregated antibody. Complement activated by the alternate pathway also enhanced attachment of trypanosomes to phagocytes, but the effect was not as rapid as it was when complement was activated by classical means.

L45 ANSWER 39 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1982:421830 CAPLUS Full-text

DOCUMENT NUMBER: 97:21830

ORIGINAL REFERENCE NO.: 97:3821a,3824a

TITLE: A monoclonal antibody defining antigenic determinants on subpopulations of mammalian neurons and *Trypanosoma cruzi* parasites

AUTHOR(S): Wood, J. N.; Hudson, L.; Jessell, T. M.; Yamamoto, M.

CORPORATE SOURCE: Dep. Immunol., St. George's Hosp. Med. Sch., London, SW17 0RE, UK

SOURCE: Nature (London, United Kingdom) (1982), 296(5852), 34-8

CODEN: NATUAS; ISSN: 0028-0836

DOCUMENT TYPE: Journal

LANGUAGE: English

AB An IgM  $\lambda$  class monoclonal antibody was raised against membranes from rat dorsal root ganglia; it defined a novel antigenic determinant expressed by subpopulations of mammalian central and peripheral neurons. In the presence of

complement it was cytotoxic to mammalian neurons in vitro. The same antibody also labeled *T. cruzi*, the protozoan responsible for Chagas' disease. Neurons labeled by the antibody were those that degenerate during this disease; thus, the labeled antigens, common to neuron and parasite, may be important in the pathogenesis of the disease.

L45 ANSWER 40 OF 42 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1982:120651 CAPLUS Full-text  
 DOCUMENT NUMBER: 96:120651  
 ORIGINAL REFERENCE NO.: 96:19791a,19794a  
 TITLE: Lectin binding sites of *Trypanosoma gambiense*  
 AUTHOR(S): Yabu, Yoshisada  
 CORPORATE SOURCE: Med. Sch., Nagoya City Univ., Nagoya, 467, Japan  
 SOURCE: Nagoya Medical Journal (1981), 26(1-2), 35-52  
 CODEN: NMJOAA; ISSN: 0027-7649  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Intact *T. gambiense* was not agglutinated by Con A, wheat germ agglutinin (WGA), soybean agglutinin (SBA), phytohemagglutinins M and P, Ricinus communis agglutinin-I, and peanut agglutinin. However, the parasites were agglutinated specifically by low concns. of these lectins after treatment by trypsin, chymotrypsin, papain, chymopapain, or pronase. *Ulex europaeus* Agglutinin-I and *Dolichos biflorus* agglutinin did not induce the agglutination of either the intact or enzyme-treated parasites. Dextranase and  $\alpha$ -amylase treatment of the protease-digested parasites did not reduce agglutination by lectins. Agglutination with lectins did not occur when an inhibitory concentration of a competitive sugar was present. Lectin-binding sites on the cell surface of *T. gambiense* were not directly associated with  $\alpha$ -1,4 or  $\alpha$ -1,6 glycosidic bonds of polysaccharides. Con A sites on the cell surface were visualized at the fine structure level with a lectin-ferritin conjugate. The electron-dense ferritin particles were distributed randomly on the enzyme-treated cell surface and flagella membrane. Con A bound at the cell surface was also visualized with horseradish peroxidase (HRP) and diaminobenzidine (DAB)-coupled reactions. A dense Con A-HRP-DAB reaction product was deposited uniformly over the entire cell surface and flagella membrane. Localization of WGA and SBA bound at the *T. gambiense* cell surface was also facilitated using HRP-DAB-coupled reactions. The fine structure distribution of these HRP-conjugated lectins was similar to that obtained with Con A-HRP-DAB prepns. Both the living and glutaraldehyde-fixed parasites gave similar agglutination results with Con A. The parasites also showed strong agglutination by Con A at low temperature. Colchicine- and cytochalasin B-pretreated parasites also showed marked agglutination by Con A. Thus, parasite agglutination by Con A differs from mammalian cell agglutination, since clustering of Con A-binding sites is not necessary for parasite agglutination.

L45 ANSWER 41 OF 42 CONFSCI COPYRIGHT 2008 CSA on STN

ACCESSION NUMBER: 2007:92194 CONFSCI  
 DOCUMENT NUMBER: 07-064010  
 TITLE: Structural Insights into Mechanisms of Dihydroorotate Oxidation and Fumarate Reduction Catalyzed by *Trypanosoma cruzi* Dihydroorotate Dehydrogenase.  
 AUTHOR: Inaoka, D. K.; Sakamoto, K.; Shimizu, H.; Shiba, T.; Kurisu, G.; Nara, T.; Aoki, T.; Kita, K.; Harada, S.  
 CORPORATE SOURCE: The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.  
 SOURCE: 000 0000: 2nd International Symposium on Diffraction Structural Biology (ISDSB 2007) (0000000). Tower Hall Funabori, Tokyo (Japan). 10-13 Sep 2007. Professor Takashi

Yamane.  
DOCUMENT TYPE: Conference  
FILE SEGMENT: DCCP  
LANGUAGE: UNAVAILABLE

L45 ANSWER 42 OF 42 WPIX COPYRIGHT 2008 THE THOMSON CORP on STN  
ACCESSION NUMBER: 2005-333264 [34] WPIX  
CROSS REFERENCE: 2005-333263  
DOC. NO. CPI: C2005-103571 [34]  
TITLE: Novel phenol derivatives, useful as antitrypanosoma  
agent for preventing and treating disease e.g.  
trypanosomiasis caused by trypanosoma  
DERWENT CLASS: B05  
INVENTOR: HOSOKAWA T; KITA K ; SAIMOTO R; SHIGEMASA Y; YABU  
Y; YAMAMOTO M; TOMOYOSHI H  
PATENT ASSIGNEE: (ARIG-N) ARIGEN INC  
COUNTRY COUNT: 107

## PATENT INFO ABBR.:

PATENT NO	KIND	DATE	WEEK	LA	PG	MAIN IPC
WO 2005037760	A1	20050428	(200534)*	JA	73	[0]
EP 1681280	A1	20060719	(200647)	EN		
AU 2004282055	A1	20050428	(200680)	EN		
JP 2005514824	X	20061228	(200702)	JA	62	
KR 2006097731	A	20060914	(200705)	KO		
US 20070208078	A1	20070906	(200759)	EN		
IN 2006DN02774	P1	20070803	(200771)	EN		

## APPLICATION DETAILS:

PATENT NO	KIND	APPLICATION	DATE
WO 2005037760	A1	WO 2004-JP15390	20041018
AU 2004282055	A1	AU 2004-282055	20041018
EP 1681280	A1	EP 2004-792559	20041018
EP 1681280	A1	WO 2004-JP15390	20041018
JP 2005514824	X	WO 2004-JP15390	20041018
KR 2006097731	A	WO 2004-JP15390	20041018
US 20070208078	A1	WO 2004-JP15390	20041018
JP 2005514824	X	JP 2005-514824	20041018
KR 2006097731	A	KR 2006-709515	20060516
US 20070208078	A1	US 2006-575653	20061213
IN 2006DN02774	P1	WO 2004-JP15390	20041018
IN 2006DN02774	P1	IN 2006-DN2774	20060517

## FILING DETAILS:

PATENT NO	KIND	PATENT NO
EP 1681280	A1 Based on	WO 2005037760 A
AU 2004282055	A1 Based on	WO 2005037760 A
JP 2005514824	X Based on	WO 2005037760 A
KR 2006097731	A Based on	WO 2005037760 A

PRIORITY APPLN. INFO: WO 2003-JP13310 20031017  
AN 2005-333264 [34] WPIX  
CR 2005-333263  
AB WO 2005037760 A1 UPAB: 20051222

NOVELTY - Phenol derivatives (I) are new.

DETAILED DESCRIPTION - Phenol derivatives of formula (I) and their salts and optical isomers are new.

X = H or halogen;

R1 = H or  $-(C_nH_{2n})-R'$ ;

n = 1-5;

R' = COOR'' (substituted by H or n carbon atoms);

R'' = H, 1-4C alkyl or  $-COR'''$ ;

R''' = pyridyl group, amino (substituted by 1-4C alkyl), phenoxyalkyl (substituted by halogen on alkyl chain or benzene ring) or phenyl group (substituted by 1-4C alkoxy or 1-4C alkoxy carbonyl group);

R2 = H or 1-4C alkyl;

R3 =  $-CHO$  or  $-COOH$ ;

R4 =  $-CH=CH-(CH_2)_p-CH_3$ ,  $-CH(OH)-(CH_2)_q-CH_3$ ,  $-CH(OH)-CH_2-CH(CH_3)-(CH_2)_2-CH=C(CH_3)_2$ ,  $-CH=CH-CH(CH_3)-(CH_2)_3-CH(CH_3)_2$ ,  $-(CH_2)_2-CH(CH_3)-(CH_2)_3-CH(CH_3)_2$  or  $-(CH_2)_8-CH_3$ ;

p = 1-12; and

q = 1-13.

INDEPENDENT CLAIMS are included for the following:

- (1) composition which contains (I);
- (2) antitrypanosoma agent which contains (I);
- (3) manufacture of antitrypanosoma agent; and
- (4) use of antitrypanosoma agent.

ACTIVITY - Protozoacide.

Antitrypanosoma effect of 3-chloro-4,6-dihydro-2-methyl-5-(3-methyl-7-(tetrahydro-5,5-dimethyl-4-oxo-2-furanyl)octyl)benzaldehyde (Ia) with respect to cyanogen resistant quinol enzyme of trypanosoma was evaluated using recombinant enzyme. (Ia) showed IC50 of 0.3 mM.

MECHANISM OF ACTION - None given.

USE - As antitrypanosoma agent for preventing and treating disease (claimed) e.g. trypanosomiasis caused by trypanosoma.

ADVANTAGE - (I) has trypanosomiasis inhibitory effect higher than ascofuranone at low concentration.

=> d his nofile

(FILE 'HOME' ENTERED AT 11:10:38 ON 20 FEB 2008)

FILE 'REGISTRY' ENTERED AT 11:10:44 ON 20 FEB 2008

```
L1          STR
L2          0 SEA SSS SAM L1
L3          20 SEA SSS FUL L1
L4          STR
L5          3 SEA SSS SAM L4
L6          46 SEA SSS FUL L4
```

FILE 'CAPLUS' ENTERED AT 11:39:00 ON 20 FEB 2008

```
L7          1 SEA ABB=ON PLU=ON L3
L8          96 SEA ABB=ON PLU=ON L6
```

FILE 'REGISTRY' ENTERED AT 11:39:18 ON 20 FEB 2008

```
L9          STR L4
L10         2 SEA SUB=L6 SSS SAM L9
L11         32 SEA SUB=L6 SSS FUL L9
```

FILE 'CAPLUS' ENTERED AT 11:42:36 ON 20 FEB 2008

```
L12         93 SEA ABB=ON PLU=ON L11
```

10/575,653

February 20, 2008

L13 ANALYZE PLU=ON L12 1-93 RN : 1785 TERMS  
D

FILE 'REGISTRY' ENTERED AT 11:43:04 ON 20 FEB 2008

L14 1 SEA ABB=ON PLU=ON 38462-04-3  
D

L15 31 SEA ABB=ON PLU=ON L11 NOT L14

FILE 'CAPLUS' ENTERED AT 11:44:59 ON 20 FEB 2008

L16 34 SEA ABB=ON PLU=ON L15

FILE 'REGISTRY' ENTERED AT 11:45:09 ON 20 FEB 2008

L17 0 SEA ABB=ON PLU=ON L15 AND C18H25CLO4/MF

L18 5 SEA ABB=ON PLU=ON L15 AND C=18  
D SCA

L19 2 SEA ABB=ON PLU=ON L18 AND H=25  
D SCA

L20 1 SEA ABB=ON PLU=ON L18 AND H=27  
D SCA

L21 0 SEA ABB=ON PLU=ON L11 AND C=21

L22 15 SEA ABB=ON PLU=ON L11 AND OC4/ES

L23 15 SEA ABB=ON PLU=ON L22 AND O=5

L24 0 SEA ABB=ON PLU=ON L23 AND N=1

L25 0 SEA ABB=ON PLU=ON H=23 AND L23

L26 3 SEA ABB=ON PLU=ON H=33 AND L23

L27 3 SEA ABB=ON PLU=ON H=31 AND L23  
D SCA L26

L28 0 SEA ABB=ON PLU=ON L26 AND C=23

L29 3 SEA ABB=ON PLU=ON L11 AND C=24  
D SCA  
E BENZALDEHYDE, 3-CHLORO-6-HYDROXY-4-METHOXY-2-METHYL-5-((2E,6E

L30 1 SEA ABB=ON PLU=ON "BENZALDEHYDE, 3-CHLORO-6-HYDROXY-4-METHOXY  
-2-METHYL-5-((2E,6E)-3-METHYL-7-((2S)-TETRAHYDRO-5,5-DIMETHYL-4  
-OXO-2-FURANYL)-2,6-OCTADIENYL)-"/CN  
D SCA  
D

L31 21 SEA ABB=ON PLU=ON L3 OR L30

FILE 'CAPLUS' ENTERED AT 12:03:19 ON 20 FEB 2008

L32 3 SEA ABB=ON PLU=ON L31

FILE 'CAPLUS, DISSABS, CONFSCI, WPIX' ENTERED AT 12:05:30 ON 20 FEB 2008  
E SAIMOTO H/AU

L33 206 SEA ABB=ON PLU=ON ("SAIMOTO H"/AU OR "SAIMOTO HIROYUKI"/AU  
OR "SAIMOTO HIRYUKI"/AU)  
E SHIGEMASA Y/AU

L34 286 SEA ABB=ON PLU=ON ("SHIGEMASA Y"/AU OR "SHIGEMASA YOSHIHIRO"/  
AU OR "SHIGEMASA YOSHIHRO"/AU)  
E KITA K/AU

L35 1742 SEA ABB=ON PLU=ON ("KITA K"/AU OR "KITA K A"/AU OR "KITA K  
F"/AU OR "KITA K K F S P"/AU OR "KITA K M B"/AU OR "KITA K M  
C"/AU OR "KITA K N G K K K K"/AU OR "KITA K S C"/AU OR "KITA  
KIYOSHI"/AU)  
E YOSHISADA Y/AU  
E YABU Y/AU

L36 85 SEA ABB=ON PLU=ON ("YABU Y"/AU OR "YABU Y T"/AU OR "YABU  
YOSHISADA"/AU)  
E HOSOKAWA T/AU

L37 1970 SEA ABB=ON PLU=ON ("HOSOKAWA TOMOYOSHI"/AU OR "HOSOKAWA  
T"/AU OR "HOSOKAWA T C O F"/AU OR "HOSOKAWA T D C"/AU OR

"HOSOKAWA T F I C L"/AU OR "HOSOKAWA T I G C L"/AU OR "HOSOKAWA  
T L"/AU OR "HOSOKAWA T N D C"/AU OR "HOSOKAWA T T G C L"/AU  
OR "HOSOKAWA T Y F L"/AU)  
E YAMAMOTO M/AU

L38 17371 SEA ABB=ON PLU=ON ("YAMAMOTO MASAICHI"/AU OR "YAMAMOTO M"/AU  
OR "YAMAMOTO M 0"/AU OR "YAMAMOTO M A"/AU OR "YAMAMOTO M B"/AU  
OR "YAMAMOTO M B K K K"/AU OR "YAMAMOTO M C"/AU OR "YAMAMOTO M  
C O M"/AU OR "YAMAMOTO M D"/AU OR "YAMAMOTO M D I L"/AU OR  
"YAMAMOTO M D K K"/AU OR "YAMAMOTO M D M"/AU OR "YAMAMOTO M D  
M C L"/AU OR "YAMAMOTO M D N P C L"/AU OR "YAMAMOTO M D P C C  
L"/AU OR "YAMAMOTO M D R L"/AU OR "YAMAMOTO M E"/AU OR  
"YAMAMOTO M E C"/AU OR "YAMAMOTO M E I"/AU OR "YAMAMOTO M  
EMILIA"/AU OR "YAMAMOTO M F"/AU OR "YAMAMOTO M F L"/AU OR  
"YAMAMOTO M F P F C L"/AU OR "YAMAMOTO M G C"/AU OR "YAMAMOTO  
M H I"/AU OR "YAMAMOTO M H L"/AU OR "YAMAMOTO M H M C L"/AU OR  
"YAMAMOTO M I"/AU OR "YAMAMOTO M I F D K K"/AU OR "YAMAMOTO M  
I P D N D I"/AU OR "YAMAMOTO M J"/AU OR "YAMAMOTO M J L"/AU OR  
"YAMAMOTO M K"/AU OR "YAMAMOTO M K C I C L"/AU OR "YAMAMOTO M  
K F"/AU OR "YAMAMOTO M K F S P"/AU OR "YAMAMOTO M K S S"/AU OR  
"YAMAMOTO M L"/AU OR "YAMAMOTO M M"/AU OR "YAMAMOTO M M C"/AU  
OR "YAMAMOTO M M C C"/AU OR "YAMAMOTO M M D K K"/AU OR  
"YAMAMOTO M M E W L"/AU OR "YAMAMOTO M M H I L"/AU OR "YAMAMOTO  
M M M"/AU OR "YAMAMOTO M M S K L"/AU OR "YAMAMOTO M M S K L P  
R C"/AU OR "YAMAMOTO M N"/AU OR "YAMAMOTO M N C I L"/AU OR  
"YAMAMOTO M N C N F"/AU OR "YAMAMOTO M N D C"/AU OR "YAMAMOTO  
M N D I"/AU OR "YAMAMOTO M N P C L"/AU OR "YAMAMOTO M O P C  
L"/AU OR "YAMAMOTO M O T"/AU OR "YAMAMOTO M P"/AU OR "YAMAMOTO  
M P I C L"/AU OR "YAMAMOTO M P R C"/AU OR "YAMAMOTO M P R C  
M"/AU OR "YAMAMOTO M R L"/AU OR "YAMAMOTO M S"/AU OR "YAMAMOTO  
M S C"/AU OR "YAMAMOTO M S E I L"/AU OR "YAMAMOTO M S K C  
L"/AU OR "YAMAMOTO M S L"/AU OR "YAMAMOTO M S L O G L"/AU OR  
"YAMAMOTO M S M I L"/AU OR "YAMAMOTO M S P C L"/AU OR "YAMAMOTO  
M S S C C L"/AU OR "YAMAMOTO M T"/AU OR "YAMAMOTO M T L"/AU  
OR "YAMAMOTO M T L T K"/AU OR "YAMAMOTO M Y"/AU OR "YAMAMOTO M  
Y F C L"/AU)

L39 21463 SEA ABB=ON PLU=ON (L33 OR L34 OR L35 OR L36 OR L37 OR L38)

L40 286 SEA ABB=ON PLU=ON L39 AND PHENOL

L41 0 SEA ABB=ON PLU=ON L40 AND ?PANASOM?

L42 0 SEA ABB=ON PLU=ON L40 AND (TRYPANOSOM? OR ?PANASOM? OR  
ANTITRYPANA?)

FILE 'CAPLUS' ENTERED AT 12:10:37 ON 20 FEB 2008  
E US2006-575653/APPS

L43 1 SEA ABB=ON PLU=ON US2006-575653/AP  
D SCA

FILE 'CAPLUS, DISSABS, CONFSCI, WPIX' ENTERED AT 12:11:33 ON 20 FEB 2008

L44 46 SEA ABB=ON PLU=ON L39 AND (?PANOSOM? OR TRYPANOS? OR  
ANTITRYPANOS?)

FILE 'CAPLUS' ENTERED AT 12:12:19 ON 20 FEB 2008  
D QUE L32  
D L32 IBIB ABS HITSTR TOT

FILE 'CAPLUS, DISSABS, CONFSCI, WPIX' ENTERED AT 12:13:05 ON 20 FEB 2008  
D QUE L44

L45 42 DUP REM L44 (4 DUPLICATES REMOVED)  
ANSWERS '1-40' FROM FILE CAPLUS  
ANSWER '41' FROM FILE CONFSCI  
ANSWER '42' FROM FILE WPIX

10/575,653

February 20, 2008

D L45 IBIB ABS TOT